



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 740 668 B1**

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
29.07.1998 Bulletin 1998/31

(21) Application number: **95907565.6**

(22) Date of filing: **19.01.1995**

(51) Int Cl.⁶: **C07D 471/14, A61K 31/395,
C07D 471/04, C07D 209/14
// (C07D 471/14, 241:00, 221:00,
209:00)**

(86) International application number:
PCT/EP95/00183

(87) International publication number:
WO 95/19978 (27.07.1995 Gazette 1995/32)

(54) TETRACYCLIC DERIVATIVES, PROCESS OF PREPARATION AND USE

TETRACYCLISCHE DERIVATE, VERFAHREN ZU IHRER HERSTELLUNG UND IHRE
VERWENDUNG

DERIVES TETRACYCLIQUES, LEURS PROCEDES DE PREPARATION ET LEUR UTILISATION

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE**
Designated Extension States:
LT SI

(30) Priority: **21.01.1994 GB 9401090**

(43) Date of publication of application:
06.11.1996 Bulletin 1996/45

(73) Proprietor: **ICOS CORPORATION**
Bothell, WA 98201 (US)

(72) Inventor: **DAUGAN, Alain Claude-Marie,**
Laboratoires Glaxo SA.
25, avenue de Québec, F-91940 Les Ulis (FR)

(74) Representative:
Gowshall, Jonathan Vallance et al
FORRESTER & BOEHMERT
Franz-Joseph-Strasse 38
80801 München (DE)

(56) References cited:
US-A- 3 917 599

- **JOURNAL OF MEDICINAL CHEMISTRY**, vol. 16, no. 5, 1973 pages 560-564, **SAXENA ET AL.** 'Agents Acting on the Central Nervous System. 15. 2-Substituted 1,2,3,4,6,7,12,12a-octahydropyrazino[2',1':6,1]pyrido[3,4-b]indoles. A New Class of Central Nervous System Depressants'
- **CHEM. PHARM. BULL.**, vol. 33, no. 8, 1985 pages 3237-3249, **ISHIDA; NAKAMURA; IRIE; OHISHI** 'A New Method for the preparation of 3,4-Dihydro- and 1,2,3,4-Tetrahydro-beta-carboline'


Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 740 668 B1

Description

This invention relates to a series of tetracyclic derivatives, to processes for their preparation, pharmaceutical compositions containing them, and their use as therapeutic agents. In particular, the invention relates to tetracyclic derivatives which are potent and selective inhibitors of cyclic guanosine 3',5'-monophosphate specific phosphodiesterase (cGMP specific PDE) having utility in a variety of therapeutic areas where such inhibition is thought to be beneficial, including the treatment of cardiovascular disorders.

Thus, according to a first aspect, the present invention provides compounds of formula (I)




and salts and solvates (e.g. hydrates) thereof, in which:

R⁰ represents hydrogen, halogen or C₁₋₆ alkyl;

R¹ represents hydrogen, C₁₋₆alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, haloC₁₋₆alkyl, C₃₋₈cydoalkyl, C₃₋₈cycloalkyl C₁₋₃alkyl, arylC₁₋₃alkyl or heteroarylC₁₋₃alkyl;


R² represents an optionally substituted monocyclic aromatic ring selected from benzene, thiophene, furan and pyridine or an optionally substituted bicyclic ring



attached to the rest of the molecule via one of the benzene ring carbon atoms and wherein the fused ring A is a 5- or 6-membered ring which may be saturated or partially or fully unsaturated and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulphur and nitrogen; and

R³ represents hydrogen or C₁₋₃ alkyl, or R¹ and R³ together represent a 3- or 4- membered alkyl or alkenyl chain.

There is further provided by the present invention a subgroup of compounds of formula (I), the subgroup comprising compounds of formula (Ia)




and salts and solvates (e.g. hydrates) thereof, in which:

R⁰ represents hydrogen, halogen or C₁₋₆ alkyl;

R¹ represents hydrogen, C₁₋₆alkyl, haloC₁₋₆alkyl, C₃₋₈cydoalkyl, C₃₋₈cycloalkylC₁₋₃alkyl, arylC₁₋₃alkyl or heteroarylC₁₋₃alkyl; and


R² represents an optionally substituted monocyclic aromatic ring selected from benzene, thiophene, furan and pyridine or an optionally substituted bicyclic ring




attached to the rest of the molecule via one of the benzene ring carbon atoms and wherein the fused ring A is a 5- or 6-membered ring which may be saturated or partially or fully unsaturated and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulphur and nitrogen.

Within R¹ above, the term "aryl" as part of an arylC₁₋₃alkyl group means phenyl or phenyl substituted by one or more (e.g. 1, 2 or 3) substituents selected from halogen, C₁₋₆alkyl, C₁₋₆alkoxy and methylenedioxy. The term "heteroaryl" as part of a heteroarylC₁₋₃alkyl group means thienyl, furyl or pyridyl each optionally substituted by one or more (e.g. 1, 2 or 3) substituents selected from halogen, C₁₋₆alkyl and C₁₋₆alkoxy. The term "C₃₋₈cydoalkyl" as a group or part of a C₃₋₈cycloalkylC₁₋₃alkyl group means a monocyclic ring comprising three to eight carbon atoms. Examples of suitable cycloalkyl rings include the C₃₋₆cydoalkyl rings cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

Within R² above, optional benzene ring substituents are selected from one or more (e.g. 1, 2 or 3) atoms or groups comprising halogen, hydroxy, C₁₋₆alkyl, C₁₋₆alkoxy, -CO₂R^b, haloC₁₋₆alkyl, haloC₁₋₆alkoxy, cyano, nitro and NR^aR^b, where R^a and R^b are each hydrogen or C₁₋₆alkyl, or R^a may also represent C₂₋₇alkanoyl or C₁₋₆alkylsulphonyl. Optional substituents for the remaining ring systems are selected from one or more (e.g. 1, 2 or 3) atoms or groups comprising halogen, C₁₋₆alkyl, C₁₋₆alkoxy and arylC₁₋₃alkyl as defined above. The bicyclic ring



may, for example, represent naphthalene, a heterocycle such as benzoxazole, benzothiazole, benzisoxazole, benzimidazole, quinoline, indole, benzothiophene or benzofuran or



(where n is an integer 1 or 2 and X and Y may each represent CH₂, O, S or NH).

In the above definitions, the term "alkyl" as a group or part of a group means a straight chain or, where available, a branched chain alkyl moiety. For example, it may represent a C₁₋₄alkyl function as represented by methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl and t-butyl. The term "alkenyl" as used herein includes straight-chained and branched alkenyl groups, such as vinyl and allyl groups. The term "alkynyl" as used herein includes straight-chained and branched alkynyl groups, suitably acetylene. The term "halogen" herein means a fluorine, chlorine, bromine or iodine atom. The term "haloC₁₋₆alkyl" means an alkyl group as defined above comprising one to six carbon atoms substituted at one or more carbon atoms by one or more (e.g. 1, 2 or 3) halogen atoms. Similarly, a haloC₁₋₆alkoxy group is a haloC₁₋₆alkyl group as defined above linked to the R² benzene ring via an oxygen atom. Examples of haloC₁₋₆alkyl groups include trifluoromethyl and 2,2,2-trifluoroethyl. An example of a haloC₁₋₆alkoxy group is trifluoromethoxy. The term "C₂₋₇alkanoyl" means a C₁₋₆alkylcarbonyl group where the C₁₋₆alkyl portion is as defined above. An example of a suitable C₂₋₇alkanoyl group is the C₂alkanoyl group acetyl.

It will be appreciated that when R³ is a halogen atom or a C₁₋₆alkyl group this substituent may be sited at any available position on the phenyl portion of the tetracyclic ring. However, a particular site of attachment is the ring 10-position.

The compounds of formula (I) may contain two or more asymmetric centres and thus can exist as enantiomers or diastereoisomers. In particular, in formula (I) above two ring chiral centres are denoted with asterisks. It is to be understood that the invention includes both mixtures and separate individual isomers of the compounds of formula (I).

The compounds of formula (I) may also exist in tautomeric forms and the invention includes both mixtures and separate individual tautomers thereof.


The pharmaceutically acceptable salts of the compounds of formula (I) which contain a basic centre are acid addition salts formed with pharmaceutically acceptable acids. Examples include the hydrochloride, hydrobromide, sulphate or bisulphate, phosphate or hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, ci-

trate, tartrate, gluconate, methanesulphonate, benzenesulphonate and p-toluenesulphonate salts. Compounds of the formula (I) can also provide pharmaceutically acceptable metal salts, in particular alkali metal salts, with bases. Examples include the sodium and potassium salts.

A particular group of compounds of the invention are those compounds of formula (I) in which R⁰ is hydrogen or halogen (e.g. fluorine), especially hydrogen.

Another particular group of compounds of the invention are those compounds of formula (I) in which R¹ represents hydrogen, C₁₋₄alkyl, haloC₁₋₄alkyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkylmethyl, pyridylC₁₋₃alkyl, furylC₁₋₃alkyl or optionally substituted benzyl. Within this particular group of compounds, examples of C₁₋₄alkyl groups are methyl, ethyl, n-propyl, i-propyl and n-butyl. Examples of C₃₋₆cycloalkylmethyl groups are cyclopropylmethyl and cyclohexylmethyl. Examples of optionally substituted, benzyl groups include benzyl and halobenzyl (e.g. fluorobenzyl).


A further particular group of compounds of the invention are those compounds of formula (I) in which R² represents an optionally substituted benzene, thiophene, furan, pvrindine or naphthalene ring or an optionally substituted bicyclic ring



(where n is 1 or 2 and X and Y are each CH₂ or O). Within this particular group of compounds, examples of substituted benzene groups are benzene substituted by one of halogen (e.g. chlorine), hydroxy, C₁₋₃alkyl (e.g. methyl, ethyl or i-propyl), C₁₋₃alkoxy (e.g. methoxy or ethoxy), -CO₂R^b, halomethyl (e.g. trifluoromethyl), halomethoxy (e.g. trifluoromethoxy), cyano, nitro or NR^aR^b where R^a and R^b are each hydrogen or methyl or R^a is acetyl; or benzene substituted by dihalo (e.g. dichloro) or by C₁₋₃alkoxy (e.g. methoxy) and one of halogen (e.g. chlorine) and hydroxy. An example of a substituted thiophene ring is a halo (e.g. bromo) substituent thiophene ring.

A still further particular group of compounds of formula I are those wherein R³ represents hydrogen or R¹ and R³ together represent a 3-membered alkyl chain.

A preferred group of compounds of the invention are the cis isomers of formula (I) represented by formula (Ib)



and mixtures thereof with their cis optical enantiomers, including racemic mixtures, and salts and solvates (e.g. hydrates) of these compounds in which R⁰ is hydrogen or halogen (e.g. fluorine), especially hydrogen and R¹, R² and R³ are as defined previously.

The single isomers represented by formula (Ib), i.e. the 6R, 12aR isomers, are particularly preferred.

Within the above definitions R¹ may preferably represent C₁₋₄alkyl (e.g. methyl, ethyl, i-propyl and n-butyl), C₃₋₆cycloalkyl (e.g. cyclopentyl) or C₃₋₆cycloalkylmethyl (e.g. cyclopropylmethyl).

R² may preferably represent a substituted benzene ring such as benzene substituted by C₁₋₃alkoxy (e.g. methoxy) or by C₁₋₃alkoxy (e.g. methoxy) and halogen (e.g. chlorine), particularly 4-methoxyphenyl or 3-chloro-4-methoxyphenyl, or R² may preferably represent 3,4-methylenedioxyphenyl.

It is to be understood that the present invention covers all appropriate combinations of particular and preferred groupings hereinabove.

Particular individual compounds of the invention include:

Cis-2,3,6,7,12,12a-hexahydro-2-(4-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thienyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methylphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-methoxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-methoxyphenyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (5aR,12R,14aS)-1,2,3,5,6,11,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1",2":4',5']pyrazino[2',1':6,1]pyrido[3,4-b]indole-5-1,4-dione;

and physiologically acceptable salts and solvates (e.g. hydrates) thereof.

A specific compound of the invention is:

(6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

and physiologically acceptable salts and solvates (e.g. hydrates) thereof.

It has been shown that compounds of the present invention are potent and selective inhibitors of cGMP specific PDE. Thus, compounds of formula (I) are of interest for use in therapy, specifically for the treatment of a variety of conditions where inhibition of cGMP specific PDE is thought to be beneficial.

As a consequence of the selective PDE V inhibition exhibited by compounds of the present invention, cGMP levels are elevated, which in turn can give rise to beneficial anti-platelet, anti-neutrophil, anti-vasospastic, vasodilatory, natriuretic and diuretic activities as well as potentiation of the effects of endothelium-derived relaxing factor (EDRF), nitro-vasodilators, atrial natriuretic factor (ANF), brain natriuretic peptide (BNP), C-type natriuretic peptide (CNP) and endothelium-dependent relaxing agents such as bradykinin, acetylcholine and 5-HT₁. The compounds of formula (I) therefore have utility in the treatment of a number of disorders, including stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, renal failure, atherosclerosis, conditions of reduced blood vessel patency (e.g. post-percutaneous transluminal coronary angioplasty), peripheral vascular disease, vascular disorders such as Raynaud's disease, inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma and diseases characterised by disorders of gut motility (e.g. irritable bowel syndrome).

It will be appreciated that references herein to treatment extend to prophylaxis as well as treatment of established conditions.

It will also be appreciated that 'a compound of formula (I)', or a physiologically acceptable salt or solvate thereof can be administered as the raw compound, or as a pharmaceutical composition containing either entity.

There is thus provided as a further aspect of the invention a compound of formula (I) for use in the treatment of stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, congestive heart failure, renal failure, atherosclerosis, conditions of reduced blood vessel patency, (e.g. post-PTCA), peripheral vascular disease, vascular disorders such as Raynaud's disease, inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility (e.g. IBS).

According to another aspect of the invention, there is provided the use of a compound of formula (I) for the manufacture of a medicament for the treatment of stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, congestive heart failure, renal failure, atherosclerosis, conditions of reduced blood vessel patency, (e.g. post-PTCA), peripheral vascular disease, vascular disorders such as Raynaud's disease, inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility (e.g. IBS).

In a further aspect, the invention provides a method of treating stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, congestive heart failure, renal failure, atherosclerosis, conditions of reduced blood vessel patency, (e.g. post-PTCA), peripheral vascular disease, vascular disorders such as Raynaud's disease, inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility (e.g. IBS) in a human or non-human animal body which comprises administering to said body a therapeutically effective amount of a compound with formula (I).

Compounds of the invention may be administered by any suitable route, for example by oral, buccal, sub-lingual,

rectal, vaginal, nasal, topical or parenteral (including intravenous, intramuscular, subcutaneous and intracoronary) administration. Oral administration is generally preferred.

For administration to man in the curative or prophylactic treatment of the disorders identified above, oral dosages of a compound of formula (I) will generally be in the range of from 0.5-800mg daily for an average adult patient (70kg). Thus for a typical adult patient, individual tablets or capsules contain from 0.2-400mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier, for administration in single or multiple doses, once or several times per day. Dosages for intravenous, buccal or sublingual administration will typically be within the range of from 0.1-400 mg per single dose as required. In practice the physician will determine the actual dosing regimen which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case but there can be individual instances in which higher or lower dosage ranges may be merited, and such are within the scope of this invention.

For human use, a compound of the formula (I) can be administered alone, but will generally be administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example, the compound may be administered orally, buccally or sublingually, in the form of tablets containing excipients such as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavouring or colouring agents. Such liquid preparations may be prepared with pharmaceutically acceptable additives such as suspending agents (e.g. methylcellulose, a semi-synthetic glyceride such as witepsol or mixtures of glycerides such as a mixture of apricot kernel oil and PEG-6 esters or mixtures of PEG-8 and caprylic/capric glycerides). A compound may also be injected parenterally, for example intravenously, intramuscularly, subcutaneously or intracoronarily. For parenteral administration, the compound is best used in the form of a sterile aqueous solution which may contain other substances, for example salts, or monosaccharides such as mannitol or glucose, to make the solution isotonic with blood.

Thus, the invention provides in a further aspect a pharmaceutical composition comprising a compound of the formula (I) together with a pharmaceutically acceptable diluent or carrier therefor.

There is further provided by the present invention a process of preparing a pharmaceutical composition comprising a compound of formula (I), which process comprises mixing a compound of formula (I) together with a pharmaceutically acceptable diluent or carrier therefor.

A compound of formula (I) may also be used in combination with other therapeutic agents which may be useful in the treatment of the above-mentioned disease states. The invention thus provides, in another aspect, a combination of a compound of formula (I) together with another therapeutically active agent.


The combination referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical compositions comprising a combination as defined above together with a pharmaceutically acceptable diluent or carrier comprise a further aspect of the invention.

The individual components of such a combination may also be administered either sequentially or simultaneously in separate pharmaceutical formulations.

Appropriate doses of known therapeutic agents for use in combination with a compound of formula (I) will be readily appreciated by those skilled in the art.


Compounds of formula (I) may be prepared by any suitable method known in the art or by the following processes which form part of the present invention. In the methods below R^0 , R^1 and R^2 are as defined in formula (I) above unless otherwise indicated.

Thus, a process (A) for preparing a compound of formula (I) wherein R^3 represents hydrogen comprises treating a compound of formula (II)



(in which Alk represents C_{1-6} alkyl, e.g. methyl or ethyl and Hal is a halogen atom, e.g. chlorine) with a primary amine R^1NH_2 in a suitable solvent such as an alcohol (e.g. methanol or ethanol) or a mixture of solvents, conveniently at a temperature of from 20°C to reflux (e.g. at about 50°C).

A compound of formula (II) may conveniently be prepared by treating a compound of formula (III)




10 with a haloacetyl halide (e.g. chloroacetyl chloride) in a suitable solvent such as a halogenated hydrocarbon (e.g. trichloromethane or dichloromethane), or an ether (e.g. tetrahydrofuran), preferably in the presence of a base such as an organic amine (e.g. a trialkylamine such as triethylamine) or an alkali metal carbonate or bicarbonate (e.g. NaHCO₃). The reaction may conveniently be effected at a temperature of from -20°C to +20°C (e.g. at about 0°C).

A compound of formula (I) may also be prepared from a compound of formula (III) in a two-step procedure via a compound of formula (II) isolated without purification.

15 Compounds of formula (I) may be prepared as individual enantiomers in two steps from the appropriate enantiomer of formula (III) or as mixtures (e.g. racemates) of either pairs of cis or trans isomers from the corresponding mixtures of either pairs of cis or trans isomers of formula (III).

Individual enantiomers of the compounds of the invention may be prepared from racemates by resolution using methods known in the art for the separation of racemic mixtures into their constituent enantiomers, for example using HPLC (high performance liquid chromatography) on a chiral column such as Hypersil naphthylurea.

20 A compound of formula (III) may conveniently be prepared from a tryptophan alkyl ester of formula (IV)



30 (where Alk is as previously defined) or a salt thereof (e.g. the hydrochloride salt) according to either of the following procedures (a) and (b). Procedure (b) is only suitable for preparing cis isomers of formula (III) and may be particularly suitable for preparing individual cis enantiomers of formula (III) from D- or L-tryptophan alkyl esters as appropriate.

35 Procedure (a)


This comprises a Pictet-Spengler cyclisation between a compound of formula (IV) and an aldehyde R²CHO. The reaction may conveniently be effected in a suitable solvent such as a halogenated hydrocarbon (e.g. dichloromethane) or an aromatic hydrocarbon (e.g. toluene) in the presence of an acid such as trifluoroacetic acid. The reaction may conveniently be carried out at a temperature of from -20°C to reflux to provide a compound of formula (III) in one step. The reaction may also be carried out in a solvent such as an aromatic hydrocarbon (e.g. benzene or toluene) under reflux, optionally using a Dean-Stark apparatus to trap the water produced.

40 The reaction provides a mixture of cis and trans isomers which may be either individual enantiomers or racemates of pairs of cis or trans isomers depending upon whether racemic or enantiomerically pure tryptophan alkyl ester was used as the starting material. Individual cis or trans enantiomers may conveniently be separated from mixtures thereof by fractional crystallisation or by chromatography (e.g. flash column chromatography) using appropriate solvents and eluents. Similarly, pairs of cis and trans isomers may be separated by chromatography (e.g. flash column chromatography) using appropriate eluents. An optically pure trans isomer may also be converted to an optically pure cis isomer using suitable epimerisation procedures. One such procedure comprises treating the trans isomer or a mixture (e.g. 1 : 1 mixture) of cis and trans isomers with methanolic or aqueous hydrogen chloride at a temperature of from 0°C to the refluxing temperature of the solution. The mixture may then be subjected to chromatography (e.g. flash column chromatography) to separate the resulting diastereoisomers, or in the procedure utilising aqueous hydrogen chloride the desired cis isomer precipitates out as the hydrochloride salt which may then be isolated by filtration.

55 Procedure (b)


This comprises a four-step procedure from a compound of formula (IV) or a salt thereof (e.g. the hydrochloride salt). The procedure is particularly suitable for preparing a 1R, 3R isomer of formula (III) from a D-tryptophan alkyl

ester of formula (IV) or a salt thereof (e.g. the hydrochloride salt). Thus, a first step (i) comprises treating a compound of formula (IV) with an acid halide R^2COHal (where Hal is as previously defined) in the presence of a base, e.g. an organic base such as a trialkylamine (for example triethylamine), to provide a compound of formula (V)




The reaction may be conveniently carried out in a suitable solvent such as a halogenated hydrocarbon (e.g. dichloromethane) or an ether (e.g. tetrahydrofuran) and at a temperature of from $-20^{\circ}C$ to $+40^{\circ}C$.

15 Step (ii) comprises treating a compound of formula (V) with an agent to convert the amide group to a thioamide group. Suitable sulfurating agents are well-known in the art. Thus, for example, the reaction may conveniently be effected by treating (V) with Lawesson's reagent. This reaction may conveniently be carried out in a suitable solvent such as an ether (e.g. dimethoxyethane) or an aromatic hydrocarbon (e.g. toluene) at an elevated temperature such as from $40^{\circ}C$ to $80^{\circ}C$ to provide a compound of formula (VI)




Step (iii) comprises treating a compound of formula (VI) with a suitable agent to provide a compound of formula (VII)



40 (where Hal is a halogen atom, e.g. iodine). The reaction may conveniently be effected by treating (VI) with an alkylating agent such as a methyl halide (e.g. methyl iodide) or an acylating agent such as an acetyl halide (e.g. acetyl chloride) in a suitable solvent such as a halogenated hydrocarbon (e.g. dichloromethane) at an elevated temperature (e.g. under reflux).


45 In step (iv) the resulting iminium halide of formula (VII) may be treated with a reducing agent such as boron hydride, e.g. sodium borohydride, to provide the desired compound of formula (III). The reduction may conveniently be effected at a low temperature, e.g. within the range of $-100^{\circ}C$ to $0^{\circ}C$, in a suitable solvent such as an alcohol (e.g. methanol).

There is further provided by the present invention a process (B) for preparing a compound of formula (I), wherein R^1 and R^3 together represent a 3- or 4-membered alkyl or alkenyl chain, which process (B) comprises cyclisation of a compound of formula (VIII)




wherein Alk represents C_{1-6} alkyl and R^1 and R^3 together represent a 3- or 4-membered chain both as hereinbefore described. The cyclisation is suitably carried out in an organic solvent or solvents, such as an alcoholic solvent (e.g. methanol) and optionally an ether solvent such as tetrahydrofuran, and in the presence of a reducing agent, aptly a palladium catalyst, such as palladium on carbon.

Conveniently a compound of formula (VIII) is prepared by reaction of a compound of formula (III) as hereinbefore described with a compound of formula (IX)



wherein Hal represents a halogen atom as hereinbefore described, R^1 and R^3 together represent a 3- or 4-membered chain as hereinbefore described and R^4 represents a protecting group, suitably a benzyloxycarbonyl group or the like. Typically the reaction is carried out in a chlorinated organic solvent, such as dichloromethane, and a tertiary amine, such as triethylamine or the like.

According to a further aspect of the present invention, there is provided a process (C) for preparing a compound of formula (I) wherein R^3 represents C_{1-3} alkyl, which process comprises cyclisation of a compound of formula (X)



wherein Alk represents C_{1-6} alkyl as hereinbefore described and R^5 represents C_{2-5} alkyl, substituted at C_1 by a halogen atom, the halogen atom being as hereinbefore described. Suitably the cyclisation is achieved by reflux for many hours, such as 22 to 26 hours, in the presence of an ether solvent, such as tetrahydrofuran, and a suitable amine as hereinafter described in the accompanying examples.

Aptly a compound of formula (X) can be prepared from a compound of formula (III) by suitable acylation techniques, such as reaction with a C_{3-6} carboxylic acid, substituted at C_2 by a halogen atom in a halogenated organic solvent, such as dichloromethane.

Compounds of formula (I) may be converted to other compounds of formula (I). Thus, for example, when R^2 is a substituted benzene ring it may be necessary or desirable to prepare the suitably substituted compound of formula (I) subsequent to process (A), (B) or (C) as above. Examples of appropriate interconversions include nitro to amino or aralkyloxy to hydroxy by suitable reducing means (e.g. using a reducing agent such as $SnCl_2$ or a palladium catalyst, such as palladium-on-carbon), or amino to substituted amino such as acylamino or sulphonylamino using standard acylating or sulphonylating conditions. In the case where R^2 represents a substituted bicyclic system, suitable interconversion can involve removal of a substituent, such as by treatment with a palladium catalyst (e.g. palladium-on-carbon) whereby, for example, a benzyl substituent may be removed from a suitable bicyclic system.

The pharmaceutically acceptable acid addition salts of the compounds of formula (I) which contain a basic centre may be prepared in a conventional manner. For example, a solution of the free base may be treated with a suitable acid, either neat or in a suitable solution, and the resulting salt isolated either by filtration or by evaporation under vacuum of the reaction solvent. Pharmaceutically acceptable base addition salts may be obtained in an analogous manner by treating a solution of a compound of formula (I) with a suitable base. Both types of salt may be formed or interconverted using ionexchange resin techniques.

Compounds of the invention may be isolated in association with solvent molecules by crystallisation from or evaporation of an appropriate solvent.

Thus, according to a further aspect of the invention, we provide a process for preparing a compound of formula (I) or a salt or solvate (e.g. hydrate) thereof which comprises process (A), (B) or (C) as hereinbefore described followed by

- i) an interconversion step; and/or either
- ii) salt formation; or
- iii) solvate (e.g. hydrate) formation.

5 There is further provided by the present invention compounds of formulae (II), (VIII), (X) and further compounds of formulae (III), (V), (VI) and (VII), with the exception for compounds (III), (V), (VI) and (VII) wherein R^o is hydrogen, R² is phenyl and Alk is methyl.

The synthesis of the compounds of the invention and of the intermediates for use therein are illustrated by the following, non-limiting Examples. In the Examples section hereinafter the following abbreviations are used: DMSO (dimethylsulphoxide), MeOH (methanol), EtOH (ethanol), DMF (dimethylformamide), EtOAc (ethyl acetate) and THF (tetrahydrofuran).

Intermediates 1 and 2

15 Methyl 1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

To a stirred solution of racemic tryptophan methyl ester (13 g) and piperonal (9.7 g) in anhydrous CH₂Cl₂ (300 mL) cooled at 0°C was added dropwise trifluoroacetic acid (9 mL) and the solution was allowed to react at ambient temperature. After 4 days, the yellow solution was diluted with CH₂Cl₂ (100 mL), washed with a saturated aqueous solution of NaHCO₃, then with water and dried over Na₂SO₄. The organic layer was evaporated to dryness under reduced pressure and the residue was purified by flash chromatography eluting with CH₂Cl₂/MeOH (99/1) to give first Intermediate 1, the cis isomer (6.5 g) m.p.: 90-93°C followed by Intermediate 2, the trans isomer (6.4 g) m.p.: 170°C.

The following compounds were obtained in a similar manner:

25 Intermediates 3 and 4

Methyl 1,2,3,4-tetrahydro-1-(4-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 4-methoxybenzaldehyde gave Intermediate 3, the cis isomer as white crystals m.p.: 142°C and Intermediate 4, the trans isomer as white crystals m.p.: 209-210°C.

Intermediate 5

35 Methyl 1,2,3,4-tetrahydro-1-(3-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer

The same method but starting from racemic tryptophan methyl ester and 3-methoxybenzaldehyde gave the title compound as white crystals m.p.: 146°C.

40 Intermediates 6 and 7

Methyl 1,2,3,4-tetrahydro-1-(4-ethoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 4-ethoxybenzaldehyde gave Intermediate 6, the cis isomer as white crystals m.p.: 180°C and Intermediate 7, the trans isomer as white crystals m.p.: 196-198°C.

Intermediates 8 and 9

50 Methyl 1,2,3,4-tetrahydro-1-(2,3-dihydrobenzo[b]furan-5-yl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 2,3-dihydrobenzo[b]furan-5-carboxaldehyde gave Intermediate 8, the cis isomer as white crystals m.p.: 106-109°C and Intermediate 9, the trans isomer as white crystals m.p.: 219-222°C.

55

Intermediates 10 and 11Methyl 1,2,3,4-tetrahydro-1-(3,4-ethylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

5 The same method but starting from racemic tryptophan methyl ester and 1,4-benzodioxan-6-carboxaldehyde gave Intermediate 10, the cis isomer as white crystals m.p.: 104-106°C and Intermediate 11, the trans isomer as white crystals m.p.: 207-209°C.

Intermediate 12

10

Methyl 1,2,3,4-tetrahydro-1-(2-chlorophenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, mixture of cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 2-chlorobenzaldehyde gave the title compound as white crystals m.p.: 154°C.

15

Intermediates 13 and 14Methyl 1,2,3,4-tetrahydro-1-(4-chlorophenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

20 The same method but starting from racemic tryptophan methyl ester and 4-chlorobenzaldehyde gave Intermediate 13, the cis isomer as white crystals m.p.: 208-209°C and Intermediate 14, the trans isomer as white crystals m.p.: 108-109°C.

Intermediates 15 and 16

25

Methyl 1,2,3,4-tetrahydro-1-(3,4-dichlorophenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 3,4-dichlorobenzaldehyde gave Intermediate 15, the cis isomer as a white solid ¹H NMR (CDCl₃) δ (ppm): 7.8-7 (m, 8H, H aromatic); 5.15 (brs, 1H, H-1); 3.9 - 3.8 (dd, 1H, H-3) 3.7 (s, 3H, CO₂CH₃); 3.2 - 3.1 (ddd, 1H, H-4) 2.9 (m, 1H, H-4); 2.4 (brs, 1H, NH) and Intermediate 16, the trans isomer as a white solid m.p.: 204°C.

30

Intermediate 17Methyl 1,2,3,4-tetrahydro-1-(1,2,3,4-tetrahydro-6-naphthyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer

The same method but starting from racemic tryptophan methyl ester and 1,2,3,4-tetrahydronaphthyl-6-carboxaldehyde gave the title compound as a white solid ¹H NMR (CDCl₃) δ (ppm): 7.7-7 (m, 8H, H aromatic); 5.2 (s, 1H, H-1); 4.0 (dd, 1H, H-3); 3.8 (s, 3H, CO₂CH₃); 3.2 (m, 1H, H-4); 3.0 (m, 1H, H-4); 2.7 (m, 4H, CH₂Ar); 1.7 (s, 4H, CH₂CH₂Ar).

40

Intermediates 18 and 19Methyl 1,2,3,4-tetrahydro-1-(2-naphthyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

45

The same method but starting from racemic tryptophan methyl ester and 2-naphthaldehyde gave Intermediate 18, the cis isomer as a white solid ¹H NMR (CDCl₃) δ (ppm): 8-6.9 (m, 12H, H aromatic); 5.4 (s, 1H, H-1); 3.95 (dd, 1H, H-3); 3.7 (s, 3H, CO₂CH₃); 3.2 (ddd, 1H, H-4); 3 (m, 1H, H-4); 2.5 (brs, 1H, NH) and Intermediate 19, the trans isomer as a white solid (0.6 g) m.p.: 119°C.

50

Intermediates 20 and 21Methyl 1,2,3,4-tetrahydro-1-(2-thienyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

55 The same method but starting from racemic tryptophan methyl ester and 2-thiophenecarboxaldehyde gave Intermediate 20, the cis isomer as a pale yellow solid m.p.: 134-137°C and Intermediate 21, the trans isomer as white crystals m.p.: 169°C.

Intermediates 22 and 23Ethyl 1,2,3,4-tetrahydro-1-(3-thienyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

5 The same method but starting from racemic tryptophan ethyl ester and 3-thiophenecarboxaldehyde gave Intermediate 22, the cis isomer as white crystals m.p.: 130°C and Intermediate 23, the trans isomer as white crystals m.p.: 182-184°C.

Intermediates 24 and 25

10

Methyl 1,2,3,4-tetrahydro-1-(5-bromo-2-thienyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 5-bromo-2-thiophenecarboxaldehyde gave Intermediate 24, the cis isomer as a cream solid m.p.: 130°C and Intermediate 25, the trans isomer as a cream solid m.p.: 205°C.

Intermediates 26 and 27

20

Methyl 1,2,3,4-tetrahydro-1-(4-bromo-2-thienyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 4-bromo-2-thiophenecarboxaldehyde gave Intermediate 26, the cis isomer as a cream solid m.p.: 200°C and Intermediate 27, the trans isomer as a cream solid m.p.: 120°C.

Intermediate 28Methyl 1,2,3,4-tetrahydro-1-(3-furyl)-9H-pyrido[3,4-b]indole-3-carboxylate, mixture of cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 3-furaldehyde gave the title compound as a yellow solid m.p.: 130°C.

Intermediates 29 and 30

35

Ethyl 1,2,3,4-tetrahydro-1-(5-methyl-2-furyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan ethyl ester and 5-methylfurfural gave Intermediate 29, the cis isomer as a oily compound ¹H NMR (CDCl₃) δ (ppm) : 7.7 (brs, 1H, NH indole); 7.5 (d, 1H, H aromatic); 7.25-6.9 (m, 3H, H aromatic); 6.15 (d, 1H, H aromatic); 5.85 (m, 1H, H aromatic); 5.25 (brs, 1H, H-1); 4.2 (q, 2H, CO₂CH₂CH₃); 3.8 (dd, 1H, H-3); 3.2-2.8 (m, 2H, H-4); 2.2 (s, 3H, CH₃); 1.25 (t, 3H, CO₂CH₂CH₃) and Intermediate 30, the trans isomer as a cream solid m.p.: 152°C.

Intermediates 31 and 32

45

Ethyl 1,2,3,4-tetrahydro-1-(4-methylphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan ethyl ester and p-tolualdehyde gave Intermediate 31, the cis isomer as white crystals m.p. : 148°C and Intermediate 32, the trans isomer as white crystals m.p.: 180°C.

Intermediates 33 and 34

50

Methyl 1,2,3,4-tetrahydro-1-(3-methylphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and m-tolualdehyde gave Intermediate 33, the cis isomer as white crystals ¹H NMR (CDCl₃) δ(ppm) : 7.6-7 (m, 9H, H aromatic); 5.2 (brs, 1H, H-1) ; 4-3.9 (dd, 1H, H-3) 3.8 (s, 3H, CO₂CH₃) ; 3.2 - 3.1 (ddd, 1H, H-4) 3 (m, 1H, H-4) ; 2.35 (s, 3H, CH₃) ; 1.7 (brs, 1H, NH) and Intermediate 34, the trans isomer as a white solid m.p.: 175°C.

Intermediates 35 and 36Methyl 1,2,3,4-tetrahydro-1-(4-trifluoromethylphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

5 The same method but starting from racemic tryptophan methyl ester and 4-trifluoromethylbenzaldehyde gave Intermediate 35, the cis isomer as pale yellow crystals m.p. : 190°C and Intermediate 36, the trans isomer as pale yellow crystals m.p.: 203°C.

Intermediates 37 and 38

10

Ethyl 1,2,3,4-tetrahydro-1-(4-cyanophenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan ethyl ester and 4-cyanobenzaldehyde gave Intermediate 37, the cis isomer as white crystals m.p. : 200°C and Intermediate 38, the trans isomer as white crystals m.p.: 156°C.

15

Intermediate 39Methyl 1,2,3,4-tetrahydro-1-(4-hydroxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer

20 The same method but starting from racemic tryptophan ethyl ester and 4-hydroxybenzaldehyde gave the title compound as pale yellow crystals ¹H NMR (DMSO) δ(ppm) : 10.3 (s, 1H, NH-indole) 9.4 (s, 1H, OH) ; 7.8 - 7.5 (m, 8H, H aromatic) ; 5.1 (brs, 1H, H-1) ; 3.9 (m, 1H, H-3) ; 3.75 (s, 3H, CO₂CH₃) 3.1 (m, 1H, H-4) ; 2.8 (m, 1H, H-4).

Intermediate 40

25

Methyl 1,2,3,4-tetrahydro-1-(3-hydroxy-4-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer

The same method but starting from racemic tryptophan methyl ester and 3-hydroxy-4-methoxybenzaldehyde gave the title compound as a yellow solid m.p. : 140-148°C.

30

Intermediate 41Methyl 1,2,3,4-tetrahydro-1-(4-hydroxy-3-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer

35 The same method but starting from racemic tryptophan methyl ester and 4-hydroxy-3-methoxybenzaldehyde gave the title compound as a cream solid m.p. : 195°C.

Intermediate 42

40

Methyl 1,2,3,4-tetrahydro-1-(4-ethylphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 4-ethylbenzaldehyde gave the cis and trans isomer of the title compound.

45 Cis isomer: white solid ¹H NMR (CDCl₃) δ(ppm) : 7.65-7.1 (m, 9H, H aromatic) ; 5.25 (brs, 1H, H-1) ; 4(dd, 1H, H-3) ; 3.9 (s, 3H, CO₂CH₃) ; 3.4 (ddd, 1H, H-4) ; 3.1 (m, 1H, H-4) ; 2.7 (q, 2H, CH₂CH₃) 1.4 (t, 3H, CH₂CH₃).
Trans isomer: white solid m.p.: 187°C.

Intermediates 43 and 44

50

Methyl 1,2,3,4-tetrahydro-1-(4-isopropylphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan ethyl ester and 4-isopropylbenzaldehyde gave Intermediate 43, the cis isomer as a white solid ¹H NMR (DMSO) δ(ppm) : 10-15 (s, 1H, NH indole) ; 7.3-6.7 (m, 8H, H aromatic) ; 5 (brs, 1H, H-1) ; 3.6 (m, 1H, H-3) ; 3.5 (s, 3H, CO₂CH₃) ; 2.95-2.5 (m, 3H, H-4 + CH-(Me)₂) 2.4 (brs, 1H, NH) ; 1(d, 6H, 2xCH₃) and Intermediate 44, the trans isomer as a white solid m.p. : 189°C.

55

Intermediates 45 and 46Ethyl 1,2,3,4-tetrahydro-1-(4-nitrophenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

5 The same method but starting from racemic tryptophan ethyl ester and 4-nitrobenzaldehyde gave Intermediate 45, the cis isomer as yellow crystals m.p. : 168°C and Intermediate 46, the trans isomer as yellow crystals m.p. : 195°C.

Intermediate 47

10 Ethyl 1,2,3,4-tetrahydro-1-(4-dimethylaminophenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, mixture of cis and trans isomers

The same method but starting from racemic tryptophan ethyl ester and 4-dimethylaminobenzaldehyde gave the title compound as white crystals m.p.: 170°C.

15

Intermediates 48 and 49Ethyl 1,2,3,4-tetrahydro-1-(3-pyridyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

20 The same method but starting from racemic tryptophan ethyl ester and 3-pyridinecarboxaldehyde gave Intermediate 48, the cis isomer as pale yellow crystals m.p. : 230-232°C and Intermediate 49, the trans isomer as white crystals m.p.: 210-214°C.

Intermediates 50 and 51

25

Methyl 1,2,3,4 tetrahydro-6-fluoro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

30 The same method but starting from racemic 5-fluoro-tryptophan methyl ester and piperonal gave Intermediate 50, the cis isomer as a cream solid m.p. : 60°C and Intermediate 51, the trans isomer as a cream solid m.p. : 213°C.

Intermediates 52 and 53Methyl 1,2,3,4-tetrahydro-6-fluoro-1-(4-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

35

The same method but starting from racemic 5-fluoro-tryptophan methyl ester and 4-methoxybenzaldehyde gave Intermediate 52, the cis isomer as a solid ¹H NMR (CDCl₃) δ (ppm) : 7.4-6.8 (m, 8H, H aromatic) ; 5.15 (brs, 1H, H-1) ; 3.9 (dd, 1H, H-3) 3.8 (s, 3H, CO₂CH₃) ; 3.2-2.9 (m, 2H, H-4) and Intermediate 53, the trans isomer as a solid m.p.: 197°C.

40

Intermediates 54 and 55(1R,3R)-Methyl 1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer and (1S,3R)-methyl 1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate trans isomer

45

To a stirred solution of D-tryptophan methyl ester (11 g) and piperonal (7.9 g) in anhydrous CH₂Cl₂ (400 mL) cooled at 0°C was added dropwise trifluoroacetic acid (7.7 mL) and the solution was allowed to react at ambient temperature. After 4 days, the yellow solution was diluted with CH₂Cl₂ (200 mL) and washed with a saturated aqueous solution of NaHCO₃, then with water (3x200 mL) and dried over Na₂SO₄. The organic layer was evaporated under reduced pressure and the residue was purified by flash chromatography eluting with dichloromethane/ethyl acetate (97/3) to give first Intermediate 54, the cis isomer (6.5 g) m.p.: 154°C followed by Intermediate 55, the trans isomer (8.4 g) m.p.: 188°C.

50

The following compounds were obtained in a similar manner :

55

Intermediate 56

(1S, 3S) Methyl-1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer
 and (1R, 3S) methyl-1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, trans
 isomer

The same method but starting from L-tryptophan methyl ester and piperonal gave the cis and trans isomers of the
 title compound.

Cis isomer: white crystals m.p.: 154°C.

Trans isomer: white crystals m.p.: 187-189°C.

Intermediates 57 and 58

(1R,3R)-Methyl 1,2,3,4-tetrahydro-1-(4-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer and

(1S,3R)-methyl 1,2,3,4-tetrahydro-1-(4-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, trans isomer

The same method but starting from D-tryptophan methyl ester and 4-methoxybenzaldehyde gave Intermediate
57, the cis isomer as white crystals m.p.: 124-125°C and Intermediate 58, trans isomer as white crystals m.p.:
 219-222°C.

Intermediates 59 and 60

(1R, 3R)-Methyl 1,2,3,4-tetrahydro-1-(3-chloro-4-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer
 and

(1S, 3R)-methyl 1,2,3,4-tetrahydro-1-(3-chloro-4-methoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, trans isomer

The same method, but starting from D-tryptophan methyl ester and 3-chloro-4-methoxybenzaldehyde gave Inter-
mediate 59, the cis isomer isolated as the hydrochloride salt as white crystals m.p.: 200°C and Intermediate 60, the
 trans isomer as white crystals m.p.: 164°C.

Intermediates 61 and 62

(1R,3R)-Methyl 1,2,3,4-tetrahydro-1-(2,3-dihydrobenzo[b]furan-5-yl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer
 and

(1S,3R)-methyl 1,2,3,4-tetrahydro-1-(5-(2,3-dihydrobenzo[b]furan))-9H-pyrido[3,4-b]indole-3-carboxylate, trans
 isomer

The same method but starting from D-tryptophan methyl ester and 2,3-dihydrobenzo[b]furan-5-carboxaldehyde
 gave Intermediate 61, the cis isomer as white crystals m.p.: 282°C and Intermediate 62, the trans isomer as white
 crystals m.p.: 204°C.

Intermediates 63 and 64

(1R,3R)-Methyl 1,2,3,4-tetrahydro-1-(5-indanyl)-9H-pyrido[3,4-b]indole-3-carboxylate cis isomer and

(1S,3R)-methyl 1,2,3,4-tetrahydro-1-(5-indanyl)-9H-pyrido[3,4-b]indole-3-carboxylate trans isomer

The same method but starting from D-tryptophan methyl ester and indan-5-carboxaldehyde gave Intermediate 63,
 the cis isomer as white crystals m.p.: 130-131°C and Intermediate 64, the trans isomer as white crystals m.p.: 196°C.

Intermediate 65

Ethyl 1,2,3,4-tetrahydro-1-(4-trifluoromethoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan ethyl ester and 4-trifluoromethoxybenzaldehyde gave cis

EP 0 740 668 B1

and trans isomers of the title compound. Cis isomer: white crystals m.p.: 88°C.
Trans isomer: white crystals m.p.: 152°C.

Intermediate 66

Methyl 1,2,3,4-tetrahydro-1-(5-methyl-2-thienyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The same method but starting from racemic tryptophan methyl ester and 5-methyl-2-thiophenecarboxaldehyde gave the cis and trans isomers of the title compound.

Cis isomer : oily compound ¹H NMR (CDCl₃) δ (ppm) : 8.4 (brs, 1H, NH-indole); 7.7 - 6.6 (m, 6H, H aromatic); 5.5 (brs, 1H, H-1); 3.9 (dd, 1H, H-3); 3.85 (s, 3H, CO₂CH₃); 3.3 - 2.9 (m, 2H, H-4); 2.5 (s, 3H, CH₃).

Trans isomer: white crystals m.p.: 194°C.

Intermediates 67 and 68

(1S,3R)-Methyl 1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate and

(1R, 3R)-methyl 1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

To a stirred solution of D-tryptophan methyl ester (obtained by treating the corresponding hydrochloride salt in water with saturated aqueous NaHCO₃ solution and extraction with CH₂Cl₂) (25.7g) and piperonal (19.4g) in anhydrous dichloromethane (700ml) cooled to 0°C was added dropwise trifluoroacetic acid (18.1ml) and the solution was allowed to react at 4°C. After 5 days, the yellow solution was diluted with dichloromethane (500ml). The organic layer was washed with a saturated aqueous solution of NaHCO₃, then with water (3 x 500ml) until the pH was neutral and dried over Na₂SO₄. The organic layer was evaporated under reduced pressure to a volume of about 500ml. The trans-isomer, which crystallised, was filtered and the filtrate was reduced to 200ml. Another fraction of the trans-isomer crystallised. The fractions of trans-isomer were combined to give the (1S,3R) isomer, Intermediate 67, as white crystals (11.4g).
mp : 188°C

[α]_D²⁰ = +32.4° (C = 1.03, CHCl₃).

The filtrate containing mainly the cis-isomer was reduced to 100ml and isopropyl ether (200ml) was added. Upon cooling, the (1R,3R) isomer, Intermediate 68, crystallised as a white solid (17.4g).

mp : 154-155°C

[α]_D²⁰ = + 24.4° (C = 1.03, CHCl₃).

Intermediate 69

(1R,3R)-Methyl 1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

Method A

Intermediate 67 (5.0g) was dissolved in methanol (150ml). Hydrogen chloride was bubbled into the solution for several minutes at 0°C and the resulting yellow solution was refluxed for 24 hours. The solvent was removed under reduced pressure and the residue was basified with a saturated aqueous solution of NaHCO₃ and extracted with dichloromethane. The organic layer was washed with water, dried over Na₂SO₄ and purified by flash chromatography eluting with dichloromethane/methanol (99/1) to give the title compound (2.3g) corresponding to an authentic sample of Intermediate 68.

Method B

Intermediate 67 (25g) was heated in 1N hydrochloric acid (78.5ml) and water (400ml) at 60°C for 36 hours. From the initial pale yellow solution, a white solid precipitated. The mixture was then allowed to cool to 0°C and the solid filtered. The solid was then washed with diisopropyl ether (3 x 200ml) and dried to give the hydrochloride salt of the title compound (20g) as a white solid.

mp (dec.): 209 - 212°C

Method C

A 1 : 1 mixture of the cis and trans isomers of Intermediates 54 and 55 (2g) was heated in 1N hydrochloric acid

(6.8ml) and water (15ml) at 50°C for 72 hours. A similar work-up as described in Method B above gave the hydrochloride salt of the title compound (1.7g) as a white solid.

Intermediate 70

(R)-N α -(3,4-Methylenedioxyphenyl)-tryptophan methyl ester

To a suspension of D-tryptophan methyl ester hydrochloride (10.2g) in anhydrous CH₂Cl₂ (150ml) cooled at 0°C was added dropwise triethylamine (12.3ml). To the resulting solution solid piperonyl chloride (8.16g) was added portionwise at the same temperature, and the mixture was stirred at room temperature for 2 h. The mixture was washed successively with water, 0.5N hydrochloric acid, water, a saturated aqueous solution of NaHCO₃ and again with water. After drying over Na₂SO₄ and evaporation of the solvent under reduced pressure, the resulting oil on trituration from hot cyclohexane afforded the title compound as a white solid (14.7g).

mp : 123-124°C

$[\alpha]_D^{20} = -844^\circ$ (c = 104, CHCl₃)

Intermediate 71

(R)-N α -(3,4-Methylenedioxyphenylthiocarbonyl)-tryptophan methyl ester

A mixture of Intermediate 70 (14g) and Lawesson's reagent (9.28g) in dimethoxyethane (280ml) was heated at 60°C under N₂ for 16 hours with stirring. The reaction mixture was evaporated to dryness and the resulting oil was dissolved in ethyl acetate, then washed successively with an aqueous saturated solution of NaHCO₃ and water and dried over Na₂SO₄. The oily residue obtained after evaporation under reduced pressure gave, on trituration from cyclohexane, a yellow powder which was filtered and washed with cooled methanol to afford the title compound (9.74g).

mp : 129-130°C

$[\alpha]_D^{20} = -186.8^\circ$ (c = 1.14, CHCl₃).

Intermediate 72

(1R,3R)-Methyl 1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

A solution of Intermediate 71 (9g) and methyl iodide (10ml) in anhydrous dichloromethane (200ml) was heated at reflux under an argon atmosphere with protection from light. After 24 hours, the solvent was removed under reduced pressure to give an orange oil which on trituration from hexane gave a solid which was washed with ether and used without further purification in the next step. This compound (13.11g) was dissolved in methanol (250ml) and the solution was cooled to -78°C. NaBH₄ (0.99g) was then added by portions and the mixture was stirred at the same temperature for 1 hour. The reaction was quenched by addition of acetone (10ml) and the solvent was removed under reduced pressure. The residue was dissolved in CH₂Cl₂, washed with water and then with brine and dried over Na₂SO₄. After evaporation of the solvent, the orange oil gave on trituration from a hot mixture of diethyl ether/cyclohexane an orange powder which was recrystallised from diethyl ether/pentane to afford the title compound as a pale yellow solid (5.15g) corresponding to an authentic sample of Intermediate 68.

Intermediate 73

(1R,3R)-Methyl 1,2,3,4-tetrahydro-2-chloroacetyl-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

Method A

To a stirred solution of Intermediate 72 (9.7g) and NaHCO₃ (2.79g) in anhydrous CHCl₃ (200ml) was added dropwise chloroacetyl chloride (5.3ml) at 0°C under N₂. The resulting mixture was stirred for 1 hour at the same temperature and diluted with CHCl₃ (100ml). Water (100ml) was then added dropwise with stirring to the mixture, followed by a saturated aqueous solution of NaHCO₃. The organic layer was washed with water until neutrality and dried over Na₂SO₄. After evaporation of the solvent under reduced pressure, the oily compound obtained was crystallised from ether to give the title compound as a pale yellow solid (9.95g).

mp : 233°C

$[\alpha]_D^{20} = -125.4^\circ$ (c = 1.17, CHCl₃).

Method B

Chloroacetyl chloride (4ml) was added dropwise to a solution of Intermediate 72 (16.1g) and triethylamine (7ml) in anhydrous CH_2Cl_2 (200ml) at 0°C under N_2 . The solution was stirred at 0°C for 30 minutes, then diluted with CH_2Cl_2 (300ml). The solution was washed with water (200ml), a saturated aqueous solution of NaHCO_3 (300ml) and brine (400ml). After drying over Na_2SO_4 and evaporation under reduced pressure, the resulting solid was washed with ether (300ml) to give the title compound as a pale yellow solid (18.3g).

Intermediate 74

Methyl 1,2,3,4-tetrahydro-6-methyl-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis and trans isomers

The cis and trans isomers of the title compound were prepared using the method described in Intermediate 1 but starting from racemic 5-methyltryptophan methyl ester and piperonal.

Cis isomer: yellow solid m.p.: 85°C .

Trans isomer: yellow solid m.p.: 185°C .

Intermediates 75 and 76

(1R, 3R)-Methyl 1,2,3,4-tetrahydro-1-(7-(4-methyl-3,4-dihydro-2H-benzo[1,4]oxazinyl))-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer and (1S, 3R)-Methyl 1,2,3,4-tetrahydro-1-(7-(4-methyl-3,4-dihydro-2H-benzo[1,4]oxazinyl))-9H-pyrido[3,4-b]indole-3-carboxylate, trans isomer

The same method, as described for intermediates 54 and 55, but starting from D-tryptophan methyl ester and 4-methyl-3,4-dihydro-2H-benzo[1,4]oxazine-7-carboxaldehyde gave Intermediate 75 the cis isomer as an oily compound ^1H NMR (CDCl_3) δ (ppm): 7.6-7.1 (m, 5H); 6.9-6.6 (m, 3H); 5.15 (br s, 1H); 4.3 (t, 2H); 4 (dd, 1H); 3.8 (s, 3H); 3.3 (t, 2H); 3.3-2.95 (m, 2H); 2.9 (s, 3H); 1.6 (br s) and intermediate 76, the trans isomer as white crystals m.p.: $119-121^\circ\text{C}$.

Intermediate 77

Methyl 1,2,3,4-tetrahydro-1-(5-(N-benzylindolinyl))-9H-pyrido[3,4-b]indole-3-carboxylate, mixture of (1R, 3R) and (1S, 3R) isomers

The same method, as described for intermediates 54 and 55, but starting from D-tryptophan methyl ester and N-benzylindoline-5-carboxaldehyde gave intermediate 77 as an oily compound.

Intermediates 78 and 79

(1R, 3R)-Methyl 1,2,3,4-tetrahydro-1-(4-carbomethoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, cis isomer and (1S, 3R)-methyl 1,2,3,4-tetrahydro-1-(4-carbomethoxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate, trans isomer

The same method, as described for intermediates 54 and 55, but starting from D-tryptophan methyl ester and methyl 4-formylbenzoate gave Intermediate 78, the cis isomer as white crystals m.p.: $157-160^\circ\text{C}$ and Intermediate 79, the trans isomer as pale yellow crystals m.p.: $124-126^\circ\text{C}$.

Intermediate 80

(1R, 3R)-Methyl 1,2,3,4-tetrahydro-2-[2-(benzyloxycarbonyl)-R-prolyl]-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

A solution of N-(benzyloxycarbonyl)-D-proline acid chloride (0.64 g, 2.4 mmol) in anhydrous dichloromethane (10 mL) was added dropwise to a stirred solution of intermediate 54 (0.7 g, 2 mmol) and triethylamine (0.33 mL, 2.4 mmol) in dichloromethane (15 mL) at -10°C . The mixture was stirred for 2 h at -10°C after which it was diluted with dichloromethane (50 mL), washed with hydrochloric acid (1N), water, a saturated solution of NaHCO_3 , a saturated NaCl solution and dried over Na_2SO_4 . Evaporation of the solvent and recrystallisation of the crude product from methanol gave the title compound as pale yellow crystals (0.75 g) m.p.: $268-270^\circ\text{C}$.

Intermediate 81

(1R, 3R)-Methyl 1,2,3,4-tetrahydro-2-[2-(benzyloxycarbonyl)-S-prolyl]-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

5

A solution of N-(benzyloxycarbonyl)-L-proline acid chloride (0.86 g, 3.2 mmol) in anhydrous dichloromethane (10 mL) was added dropwise to a stirred solution of intermediate 54 (0.91 g, 2.6 mmol) and triethylamine (0.44 mL, 3.2 mmol) in dichloromethane (20 mL) at -10°C. The mixture was stirred for 2 hours at -10°C after which it was diluted with dichloromethane (60 mL), washed with hydrochloric acid (1N), water, a saturated solution of NaHCO₃, a saturated NaCl solution and dried over Na₂SO₄. Evaporation of the solvent and recrystallisation of the crude product from methanol/water gave the title compound as pale yellow crystals (0.8 g) m.p.: 115-120°C.

10

Intermediate 82

(1R, 3R)-Methyl 1,2,3,4-tetrahydro-2-(2-chloropropionyl)-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

15

To a solution of (S)-(-)-2-chloropropionic acid (87 µl, 1 mmol) in anhydrous dichloromethane (15 mL), was added dicyclohexylcarbodiimide (0.23 g, 1.1 mmol). Intermediate 54 (0.35 g, 1 mmol) was then added and the mixture was stirred at room temperature for 20 hours. The formed precipitate of dicyclohexylurea was removed by filtration, the filtrate was evaporated in vacuo and the crude product was purified by flash chromatography eluting with toluene/ethyl acetate: 95/5. The oily compound obtained was then crystallised from ether/hexane to give the title compound as pale yellow crystals (0.31 g) m.p.: 125-127°C.

20

Intermediate 83

25

(1R, 3R)-Methyl 1,2,3,4-tetrahydro-2-(2-chloropropionyl)-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

To a solution of (R)-(+)-2-chloropropionic acid (191 µl, 2.2 mmol) in anhydrous dichloromethane (30 mL), was added dicyclohexylcarbodiimide (0.45 g, 2.2 mmol). Intermediate 54 (0.7 g, 2 mmol) was then added and the mixture was stirred at room temperature for 20 hours. The formed precipitate of dicyclohexylurea was removed by filtration, the filtrate was evaporated in vacuo and the crude product was purified by flash chromatography eluting with toluene/ethyl acetate: 95/5. The oily compound obtained was then crystallised from ether/hexane to give the title compound as pale yellow crystals (0.74 g) m.p.: 126-128°C.

30

35

Intermediates 84 and 85

(1R, 3R)-Methyl 1,2,3,4-tetrahydro-1-(3,4-dibenzyloxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate cis isomer and (1S, 3R)-methyl 1,2,3,4-tetrahydro-1-(3,4-dibenzyloxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate trans isomer

40

The same method as described for intermediates 54 and 55 but starting from D-tryptophan methyl ester and 3,4-dibenzyloxybenzaldehyde gave intermediate 84, the cis isomer as an oily compound ¹H NMR (CDCl₃) δ(ppm): 7.5 - 6.95 (m, 15H); 6.85 (s, 1H); 6.75 (s, 2H); 5.1 (s, 2H); 5 (br s, 1H); 4.95 (d, 2H) 3.85 (dd, 1H); 3.7 (s, 3H); 3.2-2.8 (m, 2H); 2.3 (br s, 1H) and intermediate 85, the trans isomer as an oily compound ¹H NMR (CDCl₃) δ (ppm) 7.6-7 (m, 15H); 6.9-6.7 (m, 3H); 5.2 (br s, 1H); 5.1 (s, 2H); 5 (s, 2H); 3.8 (t, 1H); 3.65 (s, 3H); 3.3-3 (m, 2H); 2.25 (br s, 1H).

45

Intermediate 86

50

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-dibenzyloxyphenyl)-2-methylpyrazino[2',1':6.1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from intermediate 84 and methylamine gave, after recrystallisation from dichloromethane/ether, the title compound as white crystals m.p.: 158-160°C, [α]_D²⁰ = +11.7° (c = 1.23; CHCl₃).

55

Intermediate 87

Methyl 1,2,3,4-tetrahydro-1-(5-(2-methylisoindolinyl))-9H-pyrido[3,4-b]indole-3-carboxylate, mixture of (1R,3R) and (1S,3R) isomers

The same method, as described for intermediates 54 and 55, but starting from D-tryptophan methyl ester and N-methylisoindoline-5-carboxaldehyde gave intermediate 87 as an oily compound.

Example 1

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

a) To a stirred solution of intermediate 1 (2 g) and NaHCO_3 (0.6 g) in anhydrous CHCl_3 (40 mL) was added dropwise chloroacetyl chloride (1.1 mL) at 0°C . The resulting mixture was stirred for 1 hour at the same temperature and diluted with CHCl_3 . Water (20 mL) was then added dropwise with stirring to the mixture, followed by a saturated solution of NaHCO_3 . The organic layer was washed with water until neutrality and dried over Na_2SO_4 . After evaporation of the solvent under reduced pressure, cis-methyl 1,2,3,4-tetrahydro-2-chloroacetyl-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate was obtained as an oil which was crystallised from ether (2 g, m.p.: $215\text{--}218^\circ\text{C}$) and was used without further purification in the next step.

b) To a stirred suspension of the chloroacetyl intermediate (0.34 g) in MeOH (20 mL) was added at ambient temperature a solution of methylamine (33% in EtOH) (0.37 mL) and the resulting mixture was heated at 50°C under N_2 for 14 hours. The solvent was removed under reduced pressure and the residue was dissolved in CH_2Cl_2 (50 mL). After washing with water (3x30 mL), drying over Na_2SO_4 and evaporating to dryness, the residue was purified by flash chromatography eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (99/1) and recrystallised from MeOH to give the title compound as white crystals (0.19 g) m.p.: $253\text{--}255^\circ\text{C}$.

Analysis for $\text{C}_{22}\text{H}_{19}\text{N}_3\text{O}_4$:

Calculated: C, 67.86; H, 4.92; N, 10.79;

Found: C, 67.53; H, 4.99; N, 10.62%.

The following compounds were obtained in a similar manner:

Example 2

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-10-fluoro-6-(4-methoxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from butylamine and intermediate 52 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 182°C .

Analysis for $\text{C}_{25}\text{H}_{26}\text{FN}_3\text{O}_3$ (0.1 H_2O):

Calculated: C, 68.67; H, 6.04; N, 9.61;

Found: C, 68.38; H, 6.11; N, 9.53%.

Example 3

Trans-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 2 gave, after recrystallisation from toluene, the title compound as white crystals m.p.: $301\text{--}303^\circ\text{C}$.

Analysis for $\text{C}_{22}\text{H}_{19}\text{N}_3\text{O}_4$:

Calculated: C, 67.86; H, 4.92; N, 10.79;

Found: C, 67.98; H, 4.98; N, 10.73%.

Example 4Cis-2,3,6,7,12,12a-hexahydro-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

5 The same two step procedure but starting from ammonia and intermediate 1 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 283-285°C.

Analysis for $C_{21}H_{17}N_3O_4$:

Calculated: C, 67.19; H, 4.56; N, 11.19;

Found: C, 67.04; H, 4.49; N, 11.10%.

10

Example 5Cis-2,3,6,7,12,12a-hexahydro-10-fluoro-6-(4-methoxyphenyl)-2-(2,2,2-trifluoroethyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

15

The same two step procedure but starting from 2,2,2-trifluoroethylamine and intermediate 52 gave, after recrystallisation from ethanol/diisopropyl ether, the title compound as white crystals m.p.: 190°C.

Analysis for $C_{23}H_{19}F_4N_3O_3$:

Calculated: C, 59.87; H, 4.15; N, 9.11;

20 Found: C, 59.81; H, 4.18; N, 9.21%.

Example 6Cis-2,3,6,7,12,12a-hexahydro-10-fluoro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

25

The same two step procedure but starting from methylamine and intermediate 50 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 292°C.

Analysis for $C_{22}H_{18}FN_3O_4$:

30 Calculated: C, 64.86; H, 4.45; N, 10.31;

Found: C, 64.66; H, 4.60; N, 10.21%.

Example 7(6R, 12aS)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and the trans isomer of intermediate 56 gave, after recrystallisation from toluene, the title compound as white crystals m.p.: 287-289°C.

40 Analysis for $C_{22}H_{19}N_3O_4$ (0.25 toluene):

Calculated: C, 69.16; H, 5.13; N, 10.19;

Found: C, 69.09; H, 5.14; N, 10.19%.

$[\alpha]_D^{20} = -293.4^\circ$ (C=1.28, $CHCl_3$).

45 Example 8(6S, 12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

50 The same two step procedure but starting from methylamine and intermediate 55 gave, after recrystallisation from toluene, the title compound as white crystals m.p.: 287°C.

Analysis for $C_{22}H_{19}N_3O_4$ (0.3 toluene):

Calculated: C, 69.41; H, 5.17; N, 10.08;

Found: C, 69.56; H, 5.24; N, 10.08%.

55 $[\alpha]_D^{20} = +297.9^\circ$ (C=1.21; $CHCl_3$).

Example 9

Cis-2,3,6,7,12,12a-hexahydro-2-[2-(2-pyridyl)-ethyl]6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from 2-(2-pyridyl)ethylamine and intermediate 1 gave, after recrystallisation from 2-propanol, the title compound as white crystals m.p. : 218-222°C.

Analysis for $C_{28}H_{24}N_4O_4$:

Calculated : C, 69.99 ; H, 5.03 ; N, 11.66;

Found : C, 69.92 ; H, 5.16 ; N, 11.48%.

Example 10

Cis-2,3,6,7,12,12a-hexahydro-2-(2-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from 2-pyridylmethylamine and intermediate 1 gave, after recrystallisation from DMF/water, the title compound as cream crystals m.p : 285-286°C.

Analysis for $C_{27}H_{22}N_4O_4$ (0.4 H_2O):

Calculated : C, 68.46 ; H, 4.85 ; N, 11.83;

Found : C, 68.58 ; H, 4.88 ; N, 11.90%.

Example 11

Cis-2,3,6,7,12,12a-hexahydro-2-(3-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from 3-pyridylmethylamine and intermediate 1 gave, after recrystallisation from $CH_2Cl_2/MeOH$, the title compound as cream crystals m.p.: 292-293°C.

Analysis : $C_{27}H_{22}N_4O_4$:

Calculated : C, 69.52 ; H, 4.75 ; N, 12.01;

Found : C, 69.27 ; H, 4.74 ; N, 11.37%.

Example 12

Cis-2,3,6,7,12,12a-hexahydro-2-(4-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from 4-pyridylmethylamine and intermediate 1 gave, after recrystallisation from MeOH, the title compound as pale yellow crystals m.p.: 273-274°C.

Analysis for $C_{27}H_{22}N_4O_4$ (1.8 H_2O):

Calculated : C, 65.00 ; H, 5.17 ; N, 11.23;

Found : C, 65.11 ; H, 4.85 ; N, 11.07%.

Example 13

Cis-2,3,6,7,12,12a-hexahydro-2-ethyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from ethylamine and intermediate 1 gave, after recrystallisation from methanol, the title compound as white crystals m.p. : 272-274°C.

Analysis for $C_{23}H_{21}N_3O_4$:

Calculated: C, 68.47; H, 5.25; N, 10.42;

Found: C, 68.52; H, 5.35; N, 10.53%.

Example 14

Cis-2,3,6,7,12,12a-hexahydro-2-(2,2,2-trifluoroethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

5

The same two step procedure but starting from 2,2,2-trifluoroethylamine and intermediate 1 gave, after recrystallisation from EtOH, the title compound as white crystals m.p.: 303°C.

Analysis for $C_{23}H_{18}F_3N_3O_4$:

Calculated: C,60.40;H,3.97;N,9.19;

10

Found: C,60.43;H,4.15;N,9.16%.

Example 15

Cis-2,3,6,7,12,12a-hexahydro-6-(3,4-methylenedioxyphenyl)-2-propylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

15

The same two step procedure but starting from propylamine and intermediate 1 gave, after recrystallisation from methanol, the title compound as white crystals m.p. : 270-271°C.

Analysis for $C_{24}H_{23}N_3O_4$:

Calculated: C,69.05;H,5.55;N,10.07;

20

Found: C,69.22;H,5.50;N,9.80%.

Example 16

Cis-2,3,6,7,12,12a-hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

25

The same two step procedure but starting from isopropylamine and intermediate 1 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 248-250°C.

Analysis for $C_{24}H_{23}N_3O_4$:

Calculated: C,69.05;H,5.55;N,10.07;

30

Found: C,68.86;H,5.66;N,10.21%.

Example 17

Cis-2,3,6,7,12,12a-hexahydro-2-cyclopropyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

35

The same two step procedure but starting from cyclopropylamine and intermediate 1 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 290-292°C.

Analysis for $C_{24}H_{21}N_3O_4$:

Calculated: C,69.39;H,5.10;N,10.11;

40

Found: C,69.11;H,5.20;N,9.94%.

Example 18

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

45

The same two step procedure but starting from butylamine and intermediate 1 gave, after recrystallisation from methanol/water, the title compound as white crystals m.p.: 241-243°C.

Analysis for $C_{25}H_{25}N_3O_4$:

Calculated: C,69.59;H,5.84;N,9.74;

50

Found: C,69.77;H,5.82;N,9.81 %.

55

Example 19

Trans-2,3,6,7,12,12a-hexahydro-2-butyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from butylamine and intermediate 2 gave, after recrystallisation from toluene, the title compound as white crystals m.p.: 243°C.

Analysis for C₂₅H₂₅N₃O₄:

Calculated: C,69.59;H,5.84;N,9.74;

Found: C,69.80;H,5.78;N,9.52%.

Example 20

Cis-2,3,6,7,12,12a-hexahydro-2-cyclopropylmethyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopropylmethylamine and intermediate 1 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 217-218°C.

Analysis for C₂₅H₂₅N₃O₄:

Calculated: C,69.92;H,5.40;N,9.78;

Found: C,70.02;H,5.47;N,9.84%.

Example 21

Cis-2,3,6,7,12,12a-hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopentylamine and intermediate 1 gave, after recrystallisation from acetone, the title compound as white crystals m.p.: 270°C.

Analysis for C₂₆H₂₅N₃O₄:

Calculated: C,70.41;H,5.68;N,9.47;

Found: C,70.58; H,5.63; N,9.38%.

Example 22

Cis-2,3,6,7,12,12a-hexahydro-2-cyclohexyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclohexylamine and intermediate 1 gave, after recrystallisation from methanol/water, the title compound as white crystals m.p.: 268-269°C.

Analysis for C₂₇H₂₇N₃O₄:

Calculated: C,70.88;H,5.95;N,9.18;

Found: C,70.82;H,5.89;N,9.21%.

Example 23

Cis-2,3,6,7,12,12a-hexahydro-2-benzyl-6-(3,4-methylenedioxyphenyl) pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from benzylamine and intermediate 1 gave, after recrystallisation from dichloromethane/hexane, the title compound as white crystals m.p.: 285-287°C.

Analysis for C₂₈H₂₃N₃O₄(1 H₂O):

Calculated: C,69.55;H,5.21;N,8.69;

Found: C,69.30;H,5.06;N,8.48%.

Example 24

Cis-2,3,6,7,12,12a-hexahydro-2-(4-fluorobenzyl)-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from 4-fluorobenzylamine and intermediate 1 gave, after recrystallisation from acetone, the title compound as white crystals m.p.: 281-283°C.

Analysis for $C_{28}H_{22}FN_3O_4$:

Calculated: C,69.56;H,4.59;F,3.93;N,8.69;

Found: C,69.54;H,4.58; F,3.82;N,8.63%.

Example 25

Cis-2,3,6,7,12,12a-hexahydro-6-(4-methoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 3 gave, after recrystallisation from 2-propanol, the title compound as white crystals m.p.: 257-263°C.

Analysis for $C_{22}H_{21}N_3O_3$:

Calculated: C,70.38;H,5.64;N,11.19;

Found: C,70.11;H,5.55;N,11.15%.

Example 26

Trans-2,3,6,7,12,12a-hexahydro-6-(4-methoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 4 gave, after recrystallisation from diisopropyl ether, the title compound as white crystals m.p.: 225-228°C.

Analysis for $C_{22}H_{21}N_3O_3$:

Calculated: C,70.38;H,5.64;N,11.19;

Found: C,70.34;H,5.77;N,11.19%.

Example 27

Cis-2,3,6,7,12,12a-hexahydro-2-ethyl-6-(4-methoxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from ethylamine and intermediate 3 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 245-255°C.

Analysis for $C_{23}H_{23}N_3O_3$:

Calculated: C,70.93;H,5.95;N,10.79;

Found: C,70.74;H,6.06;N,10.87%.

Example 28

Cis-2,3,6,7,12,12a-hexahydro-6-(4-methoxyphenyl)-2-(2,2,2-trifluoroethyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from 2,2,2-trifluoroethylamine and intermediate 3 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 232°C.

Analysis for $C_{23}H_{20}F_3N_3O_3$:

Calculated: C,62.30;H,4.55;N,9.48;

Found: C,62.08;H,4.66;N,9.54%.

Example 29

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methoxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from butylamine and intermediate 3 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 157°C.

Analysis for $C_{25}H_{27}N_3O_3(0.5H_2O)$:
 Calculated: C, 70.40; H, 6.62; N, 9.85;
 Found: C, 70.25; H, 6.60; N, 9.83%.

5 Example 30

Trans-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methoxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

10 The same two step procedure but starting from butylamine and intermediate 4 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 212-214°C.

Analysis for $C_{25}H_{27}N_3O_3$:
 Calculated: C, 71.92; H, 6.52; N, 10.06;
 Found: C, 71.81; H, 6.55; N, 10.03%.

15 Example 31

Cis-2,3,6,7,12,12a-hexahydro-6-(4-methoxyphenyl)-2-cyclopropylmethylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

20 The same two step procedure but starting from cyclopropylmethylamine and intermediate 3 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 180-185°C.

Analysis for $C_{25}H_{25}N_3O_3(0.5H_2O)$:
 Calculated: C, 70.74; H, 6.17; N, 9.90;
 Found: C, 70.91; H, 6.16; N, 9.80%.

25

Example 32

Cis-2,3,6,7,12,12a-hexahydro-2-benzyl-6-(4-methoxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

30 The same two step procedure but starting from benzylamine and intermediate 3 gave, after recrystallisation from acetone, the title compound as white crystals m.p.: 275-279°C.

Analysis for $C_{28}H_{25}N_3O_3$:
 Calculated: C, 74.48; H, 5.58; N, 9.31;
 Found: C, 74.53; H, 5.60; N, 9.20%.

35

Example 33

Cis-2,3,6,7,12,12a-hexahydro-6-(3-methoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

40 The same two step procedure but starting from methylamine and intermediate 5 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 267-269°C.

Analysis for $C_{22}H_{21}N_3O_3$:
 Calculated: C, 70.38; H, 5.64; N, 11.19;
 Found: C, 70.32; H, 5.59; N, 11.25%.

45

Example 34

Cis-2,3,6,7,12,12a-hexahydro-6-(4-ethoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

50 The same two step procedure but starting from methylamine and intermediate 6 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 247-248°C.

Analysis for $C_{23}H_{23}N_3O_3$:
 Calculated: C, 70.93; H, 5.95; N, 10.79;
 Found: C, 71.23; H, 5.95; N, 10.63%.

55

Example 35

Cis-2,3,6,7,12,12a-hexahydro-6-(4-ethoxyphenyl)-2-cyclopropylmethylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopropylmethylamine and intermediate 6 gave, after recrystallisation from 2-propanol, the title compound as white crystals m.p.: 160-162°C.

Analysis for $C_{26}H_{27}N_3O_3$:

Calculated: C, 72.71; H, 6.34; N, 9.78;

Found: C, 72.28; H, 6.39; N, 9.71%.

Example 36

Cis-2,3,6,7,12,12a-hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 8 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 292-294°C.

Analysis for $C_{23}H_{21}N_3O_3$:

Calculated: C, 71.30; H, 5.46; N, 10.85;

Found: C, 71.15; H, 5.56; N, 10.84%.

Example 37

Cis-2,3,6,7,12,12a-hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-cyclopropylmethyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopropylmethylamine and intermediate 8 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 165-166°C.

Analysis for $C_{26}H_{25}N_3O_3$:

Calculated: C, 73.05; H, 5.89; N, 9.83;

Found: C, 73.08; H, 5.97; N, 9.87%.

Example 38

Cis-2,3,6,7,12,12a-hexahydro-6-(3,4-ethylenedioxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 10 gave, after recrystallisation from acetone, the title compound as white crystals m.p.: 303-305°C.

Analysis for $C_{23}H_{21}N_3O_4$:

Calculated: C, 68.47; H, 5.25; N, 10.42;

Found: C, 68.35; H, 5.31; N, 10.27%.

Example 39

Cis-2,3,6,7,12,12a-hexahydro-6-(3,4-ethylenedioxyphenyl)-2-cyclopropylmethylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopropylmethylamine and intermediate 10 gave, after recrystallisation from dichloromethane/ether, the title compound as white crystals m.p.: 288-290°C.

Analysis for $C_{26}H_{25}N_3O_4$:

Calculated: C, 70.41; H, 5.68; N, 9.47;

Found: C, 70.15; H, 5.62; N, 9.30%.

Example 40Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(2-chlorophenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

5 The same two step procedure but starting from butylamine and intermediate 12 gave, after recrystallisation from methanol/water, the title compound as white crystals m.p.: 146°C.

Analysis for $C_{24}H_{24}ClN_3O_2(0.75 H_2O)$:

Calculated: C,66.20;H,5.90;N,9.65;

Found: C,66.15;H,5.95;N,9.69%.

10

Example 41Cis-2,3,6,7,12,12a-hexahydro-6-(4-chlorophenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

15 The same two step procedure but starting from methylamine and intermediate 13 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 274°C.

Analysis for $C_{21}H_{18}ClN_3O_2(0.25 H_2O)$:

Calculated: C,65.63;H,4.85;N,10.93;

Found: C,65.39;H,4.84;N,10.85%.

20

Example 42Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-chlorophenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

25 The same two step procedure but starting from butylamine and intermediate 13 gave, after recrystallisation from ethanol/water, the title compound as white crystals m.p.: 164-166°C.

Analysis for $C_{24}H_{24}ClN_3O_2$:

Calculated: C,68.32;H,5.73;Cl,8.40;N,9.96;

Found: C,68.48;H,5.64;Cl,8.37;N,9.99%.

30

Example 43Cis-2,3,6,7,12,12a-hexahydro-6-(3,4-dichlorophenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

35 The same two step procedure but starting from methylamine and intermediate 15 gave, after recrystallisation from ethanol/DMF, the title compound as white crystals m.p.: >260°C.

Analysis for $C_{21}H_{17}Cl_2N_3O_2(0.5 H_2O)$:

Calculated: C,59.39;H,4.29;N,9.93;

Found: C,59.32;H,4.16;N,9.99%.

40

Example 44Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-phenylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

45 The same two step procedure but starting from butylamine and cis-methyl 1,2,3,4-tetrahydro-1-phenyl-9H-pyrido[3,4-b]indole-3-carboxylate¹ gave, after recrystallisation from methanol/water, the title compound as white crystals m.p.: 243-245°C.

Analysis for $C_{24}H_{25}N_3O_2$:

Calculated: C,74.39;H,6.50;N,10.84;

50

Found: C,74.54;H,6.51;N,10.86%.

1. D. Soerens et al., J. Org. Chem. 44, 535 - 545 (1979).

Example 45

55 Cis-2,3,6,7,12,12a-hexahydro-2-benzyl-6-phenylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from benzylamine and cis-methyl 1,2,3,4-tetrahydro-1-phenyl-9H-pyrido[3,4-b]indole-3-carboxylate gave, after recrystallisation from methanol, the title compound as white crystals m.p.:

193-195°C.

Analysis for $C_{27}H_{23}N_3O_2$:

Calculated: C, 76.94; H, 5.50; N, 9.97;

Found: C, 77.23; H, 5.54; N, 9.97%.

5

Example 46

Trans-2,3,6,7,12,12a-hexahydro-2-benzyl-6-phenyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

10 The same two step procedure but starting from benzylamine and cis-methyl-1,2,3,4-tetrahydro-1-phenyl-9H-pyrido[3,4-b]indole-3-carboxylate gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 284°C.

Analysis for $C_{27}H_{23}N_3O_2$:

Calculated: C, 76.94; H, 5.50; N, 9.97;

15 Found: C, 76.88; H, 5.45; N, 9.89%.

Example 47

20 Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(1,2,3,4-tetrahydro-6-naphthyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 17 gave, after recrystallisation from methanol, the title compound as white crystals m. p. : >260°C.

Analysis for $C_{25}H_{25}N_3O_2$:

25 Calculated: C, 75.16; H, 6.31; N, 10.52;

Found: C, 74.93; H, 6.43; N, 10.63%.

Example 48

30 Cis-2,3,6,7,12,12a-hexahydro-2-isopropyl-6-(1,2,3,4-tetrahydro-6-naphthyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from isopropylamine and intermediate 17 gave, after recrystallisation from the title compound as off-white crystals m.p.: 244-246°C.

35 Analysis for $C_{27}H_{29}N_3O_2$ (0.25 H_2O):

Calculated: C, 75.06; H, 6.88; N, 9.73;

Found: C, 75.00; H, 6.83; N, 9.69%.

Example 49

40

Cis-2,3,6,7,12,12a-hexahydro-2-cyclopropylmethyl-6-(1,2,3,4-tetrahydro-6-naphthyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopropylmethylamine and intermediate 17 gave, after recrystallisation from ethanol/pentane, the title compound as white crystals m.p. : 125°C.

45 Analysis for $C_{28}H_{29}N_3O_2$ (0.25 H_2O):

Calculated: C, 75.73; H, 6.70; N, 9.46;

Found: C, 75.45; H, 6.86; N, 9.14%.

50

Example 50

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(2-naphthyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 18 gave, after recrystallisation from dichloromethane/methanol, the title compound as white crystals m.p. : >260°C.

55

Analysis for $C_{25}H_{21}N_3O_2$ (0.25 H_2O):

Calculated: C, 75.08; H, 5.42; N, 10.51;

Found: C, 75.35; H, 5.42; N, 10.49%.

Example 51Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(2-thienyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

5 The same two step procedure but starting from butylamine and intermediate 20 gave, after recrystallisation from ethanol, the title compound as white crystals m.p. : 226°C.

Analysis for $C_{22}H_{23}N_3O_2S$:

Calculated: C,67.15;H,5.89;N,10.68;

Found:C,67.39;H,5.88;N,10.77%.

10

Example 52Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thienyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

15 The same two step procedure but starting from methylamine and intermediate 24 gave, after recrystallisation from ethanol, the title compound as a cream powder m.p. : 258°C.

Analysis for $C_{19}H_{16}BrN_3O_2S$:

Calculated: C,53.03;H,3.75;N,9.76;

Found:C,53.01;H,3.78;N,9.69%.

20

Example 53Cis-2,3,6,7,12,12a-hexahydro-6-(4-bromo-2-thienyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

25 The same two step procedure but starting from methylamine and intermediate 26 gave, after recrystallisation from ethanol, the title compound as white crystals mp.: 292°C.

Analysis for $C_{19}H_{16}BrN_3O_2S$ (0.25H₂O):

Calculated: C,52.48;H,3.82;N,9.66;

Found:C,52.46;H,3.81 ;N,9.60%.

30

Example 54Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thienyl)-2-cyclopropylmethylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

35

The same two step procedure but starting from cyclopropylmethylamine and intermediate 24 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 190°C.

Analysis for $C_{22}H_{20}BrN_3O_2S$:

Calculated: C,56.18;H,4.29;N,8.93;

Found:C,55.92;H,4.28;N,8.74%.

40

Example 55Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thienyl)-2-cyclopentylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

45

The same two step procedure but starting from cyclopentylamine and intermediate 24 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 252°C.

Analysis for $C_{23}H_{22}BrN_3O_2S$:

Calculated: C,57.03;H,4.58;N,8.67;

Found:C,56.87;H,4.66;N,8.68%.

50

Example 56Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(5-methyl-2-thienyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

55

The same two step procedure but starting from methylamine and the cis isomer of intermediate 66 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 282°C.

Analysis for $C_{20}H_{19}N_3O_2S$ (0.25H₂O):

Calculated: C,64.93;H,5.31;N,11.36;
Found:C,64.84;H,5.28;N,11.08%.

Example 57

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3-thienyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 22 gave, after recrystallisation from acetone, the title compound as white crystals m.p.: 290-295°C.

Analysis for $C_{19}H_{17}N_3O_2S$:
Calculated: C,64.94;H,4.88;N,11.96;
Found: C, 64.81 ; H,4.95 ; N,11.68%.

Example 58

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(3-thienyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from butylamine and intermediate 22 gave, after recrystallisation from methanol, the title compound as white crystals m.p. : 236-239°C.

Analysis for $C_{22}H_{23}N_3O_2S$:
Calculated: C,67.15;H,5.89;N,10.68;S,8.15;
Found:C,67.42;H,5.76;N,10.57;S,8.01 %.

Example 59

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3-furyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and the cis isomer of intermediate 28 gave, after recrystallisation from ether, the title compound as a white solid m.p. : 250°C.

Analysis for $C_{19}H_{17}N_3O_3$ (0.5H₂O):
Calculated: C,66.27;H,5.27;N,12.20;
Found:C,66.33;H,5.48;N,12.02%.

Example 60

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(5-methyl-2-furyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 29 gave, after recrystallisation from ethanol, the title compound as a cream powder m.p.: 303°C.

Analysis for $C_{20}H_{19}N_3O_3$ (0.25H₂O):
Calculated: C,67.88;H,5.55;N,11.87;
Found:C,67.90;H,5.50;N,11.98%.

Example 61

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 31 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: >260°C.

Analysis for $C_{22}H_{21}N_3O_2$ (0.25 H₂O):
Calculated: C,72.61;H,5.95;N,11.55;
Found:C,72.73;H,5.96;N,11.59%.

Example 62

Cis-2,3,6,7,12,12a-hexahydro-2-isopropyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from isopropylamine and intermediate 31 gave, after recrystallisation

from the title compound as white crystals m.p.: 170°C.

Analysis for $C_{24}H_{25}N_3O_2$ (0.5H₂O):
 Calculated: C,72.70;H,6.61;N,10.60;
 Found: C,73.06;H,6.43;N,9.66%.

5

Example 63

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

10 The same two step procedure but starting from butylamine and intermediate 31 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 194°C.

Analysis for $C_{25}H_{27}N_3O_2$ (0.5H₂O):
 Calculated: C,73.15;H,6.87;N,10.24;
 Found: C,73.01;H,6.84;N,10.26%.

15

Example 64

Cis-2,3,6,7,12,12a-hexahydro-2-cyclopropylmethyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dione

20

The same two step procedure but starting from cyclopropylmethylamine and intermediate 31 gave, after recrystallisation from methanol/water, the title compound as white crystals m.p.: 194°C.

Analysis for $C_{25}H_{25}N_3O_2$ (1.1 H₂O):
 Calculated: C,71.61;H,6.54;N,10.02;
 Found: C,71.42;H,6.07;N,9.95%.

25

Example 65

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

30

The same two step procedure but starting from methylamine and intermediate 33 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: >260°C.

Analysis for $C_{22}H_{21}N_3O_2$:
 Calculated: C,73.52;H,5.89;N,11.69;
 Found: C,73.60;H,5.97;N, 11.66%.

35

Example 66

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-trifluoromethylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

40

The same two step procedure but starting from butylamine and intermediate 35 gave, after recrystallisation from methanol/water, the title compound as white crystals m.p.: 155°C.

Analysis for $C_{25}H_{24}F_3N_3O_2$ (0.5H₂O):
 Calculated: C,64.65;H,5.43;N,9.05;
 Found: C,64.78;H,5.40;N,9.01 %.

45

Example 67

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(4-trifluoromethoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

50

The same two step procedure but starting from methylamine and the cis isomer of intermediate 65 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 174-180°C.

Analysis for $C_{22}H_{18}F_3N_3O_3$ (0.5H₂O):
 Calculated: C,60.27;H,4.37;N,9.58;
 Found: C,60.24;H,4.28;N,9.50%.

55

Example 68Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(4-hydroxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

5 The same two step procedure but starting from methylamine and intermediate 39 gave, after recrystallisation from methanol, the title compound as yellow crystals m.p.: 179-180°C.

Analysis for $C_{21}H_{19}N_3O_3(1.25H_2O)$:

Calculated: C, 65.70; H, 5.64; N, 10.94;

Found: C, 65.46; H, 5.45; N, 10.92%.

Example 69Cis-2,3,6,7,12,12a-hexahydro-6-(3-hydroxy-4-methoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

15 The same two step procedure but starting from methylamine and intermediate 40 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 320°C.

Analysis for $C_{22}H_{21}N_3O_4(0.25H_2O)$:

Calculated: C, 66.74; H, 5.47; N, 10.61;

20 Found: C, 66.72; H, 5.46; N, 10.53%.

Example 70Cis-2,3,6,7,12,12a-hexahydro-6-(4-hydroxy-3-methoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

25 The same two step procedure but starting from methylamine and intermediate 41 gave, after recrystallisation from dichloromethane/ethanol, the title compound as yellow crystals m.p.: 264-265°C.

Analysis for $C_{22}H_{21}N_3O_4$:

30 Calculated: C, 67.51; H, 5.41; N, 10.74;

Found: C, 67.05; H, 5.41; N, 10.62%.

Example 71Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-cyanophenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

35 The same two step procedure but starting from butylamine and intermediate 37 gave, after recrystallisation from methanol/water, the title compound as white crystals m.p.: 246°C.

Analysis for $C_{25}H_{24}N_4O_2(1H_2O)$:

40 Calculated: C, 69.75; H, 6.09; N, 13.01;

Found: C, 69.50; H, 5.96; N, 12.86%.

Example 72Cis-2,3,6,7,12,12a-hexahydro-6-(4-ethylphenyl)-2-isopropylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

45 The same two step procedure but starting from isopropylamine and the cis isomer of intermediate 42 gave, after recrystallisation from n-pentane, the title compound as white crystals m.p.: 130°C.

Analysis for $C_{26}H_{27}N_3O_2(0.5H_2O)$:

50 Calculated: C, 73.15; H, 6.87; N, 10.24;

Found: C, 73.39; H, 7.08; N, 9.81%.

Example 73Cis-2,3,6,7,12,12a-hexahydro-6-(4-ethylphenyl)-2-cyclopropylmethylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

55 The same two step procedure but starting from cyclopropylmethylamine and the cis isomer of intermediate 42 gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 160°C.

Analysis for $C_{26}H_{27}N_3O_2$:
 Calculated: C, 75.52; H, 6.58; N, 10.16;
 Found: C, 75.54; H, 6.62; N, 10.08%.

5 Example 74

Cis-2,3,6,7,12,12a-hexahydro-6-(4-isopropylphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 43 gave, after recrystallisation from
 10 ethanol, the title compound as white crystals m.p.: 244°C.

Analysis for $C_{24}H_{25}N_3O_2$:
 Calculated: C, 74.39; H, 6.50; N, 10.84;
 Found: C, 74.27; H, 6.53; N, 11.05%.

15 Example 75

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-nitrophenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from butylamine and intermediate 45 gave, after recrystallisation from
 20 methanol, the title compound as white crystals m.p.: 182°C.

Analysis for $C_{24}H_{24}N_4O_4$ ($0.25H_2O$):
 Calculated: C, 65.97; H, 5.65; N, 12.82;
 Found: C, 65.92; H, 5.62; N, 12.96%.

25 Example 76

Cis-2,3,6,7,12,12a-hexahydro-6-(4-dimethylaminophenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and the cis isomer of intermediate 47 gave after
 30 recrystallisation from methanol, the title compound as white crystals m.p.: 266°C.

Analysis for $C_{23}H_{24}N_4O_2$:
 Calculated: C, 71.11; H, 6.23; N, 14.42;
 Found: C, 71.19; H, 6.24; N, 14.34%.

35 Example 77

Cis-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3-pyridyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 48 gave after recrystallisation from
 40 chloroform, the title compound as white crystals m.p.: 312°C.

Analysis for $C_{20}H_{18}N_4O_2$:
 Calculated: C, 69.35; H, 5.24; N, 16.17;
 Found: C, 69.08; H, 5.20; N, 16.19%.

45 Example 78

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

50 a) To a stirred solution of intermediate 54 (0.5 g) and $NaHCO_3$ (0.14 g) in anhydrous $CHCl_3$ (20 mL) was added dropwise chloroacetyl chloride (0.27 mL) at 0°C. The resulting mixture was stirred for 1 hour at the same temperature and diluted with $CHCl_3$ (20 mL). Water (10 mL) was then added dropwise with stirring to the mixture, followed by a saturated solution of $NaHCO_3$. The organic layer was washed with water until neutrality and dried over Na_2SO_4 .
 55 After evaporation of the solvent under reduced pressure, (6R,12aR)-methyl 1,2,3,4-tetrahydro-2-chloroacetyl-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate was obtained as an oil which was crystallised from ether to give a solid (0.38 g, m.p.: 233°C) which was used without further purification in the next step.

b) To a stirred suspension of the chloroacetyl intermediate (0.37 g) in MeOH (20 mL) was added at room temper-

EP 0 740 668 B1

ature a solution of methylamine (33% in EtOH) (0.4 mL) and the resulting mixture was heated at 50°C under N₂ for 16 hours. The solvent was removed under reduced pressure and the residue was dissolved in CH₂Cl₂ (50 mL). After washing with water (3x20 mL), drying over Na₂SO₄ and evaporating to dryness, the residue was purified by flash chromatography eluting with CH₂Cl₂/MeOH (99/1) and recrystallised from 2-propanol to give the title compound as white crystals (0.22 g) m.p.: 302-303°C.

Analysis for C₂₂H₁₉N₃O₄:

Calculated: C, 67.86; H, 4.92; N, 10.79;

Found: C, 67.77; H, 4.92; N, 10.74%.

20° [α]_D²⁰ = +71.0° (C=1.00; CHCl₃).

The following compounds were obtained in a similar manner:

Example 79

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from isopropylamine and intermediate 54 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 290-293°C.

Analysis for C₂₄H₂₃N₃O₄:

Calculated: C, 69.05; H, 5.55; N, 10.07;

Found: C, 69.06; H, 5.49; N, 10.12%.

[α]_D²⁰ = +52.6° (C=1.14; CHCl₃).

Example 80

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-butyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from butylamine and intermediate 54 gave, after recrystallisation from toluene/hexane, the title compound as white crystals m.p.: 209-210°C.

Analysis for C₂₅H₂₅N₃O₄:

Calculated: C, 69.59; H, 5.84; N, 9.74;

Found: C, 69.70; H, 5.93; N, 9.74%.

[α]_D²⁰ = +50.2° (C=0.53; CHCl₃).

Example 81

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isobutyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from isobutylamine and intermediate 54 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 227-228°C.

Analysis for C₂₅H₂₅N₃O₄:

Calculated: C, 69.59; H, 5.84; N, 9.74;

Found: C, 69.52; H, 5.87; N, 9.74%.

[α]_D²⁰ = +45° (C=1.04; CHCl₃).

Example 82

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopentylamine and intermediate 54 gave, after recrystallisation from ether, the title compound as white crystals m.p.: 237-239°C.

Analysis for C₂₆H₂₅N₃O₄:

Calculated: C, 70.41; H, 5.68; N, 9.47;

Found: C, 70.13; H, 5.67; N, 9.42%.

$[\alpha]_D^{20} = +36.6^\circ$ (C=0.98; CHCl₃).

Example 83

5 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-2-cyclohexylmethyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclohexylmethylamine and the cis isomer of intermediate 56 gave, after recrystallisation from 2-propanol the title compound as white crystals m.p.: 209°C.

10 Analysis for C₂₈H₂₉N₃O₄:
Calculated: C,71.32;H,6.20;N,8.91;
Found:C,71.30;H,6.29;N,8.74%.
 $[\alpha]_D^{20} = +40.0^\circ$ (C=0.99; CHCl₃).

15 Example 84

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

20 The same two step procedure but starting from cyclopropylmethylamine and intermediate 57 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 204-205°C.

Analysis for C₂₅H₂₅N₃O₃(0.5H₂O):
Calculated: C,70.74;H,6.17;N,9.90;
Found:C,70.98;H,6.09;N,9.92%.
25 $[\alpha]_D^{20} = +54.1^\circ$ (C=1.03; CHCl₃).

Example 85

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-butyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

30 The same two step procedure but starting from butylamine and intermediate 57 gave, after recrystallisation from 2-propanol, the title compound as white crystals m.p.: 183-184°C.

Analysis for C₂₅H₂₇N₃O₃(0.5H₂O):
Calculated: C,70.40;H,6.62;N,9.85;
35 Found:C,70.55;H,6.64;N,9.92%.
 $[\alpha]_D^{20} = +45.4^\circ$ (C=1.04; CHCl₃).

Example 86

40 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopentylamine and intermediate 57 gave, after recrystallisation from ether, the title compound as white crystals m.p.: 210-211°C.

45 Analysis for C₂₆H₂₇N₃O₃:
Calculated: C,72.71;H,6.34;N,9.78;
Found:C,72.53;H,6.39;N,9.53%.
 $[\alpha]_D^{20} = +29.8^\circ$ (C=1.07; CHCl₃).

50 Example 87

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-methoxyphenyl)-2-cyclopropylmethyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

55 The same two step procedure but starting from cyclopropylmethylamine and intermediate 59 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 218-219°C.

Analysis for C₂₅H₂₄ClN₃O₃ (0.25 H₂O):
Calculated: C,66.08;H,5.43;N,9.25; Cl, 7.80;

Found: C, 66.11; H, 5.33; N, 9.03; Cl, 7.74%.

$[\alpha]_D^{20} = +49.4^\circ$ (C=1.03; CHCl₃).

Example 88

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3-chloro-4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from cyclopentylamine and intermediate 59 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 260-262°C.

Analysis for C₂₆H₂₆ClN₃O₃:

Calculated: C, 67.31; H, 5.65; Cl, 7.64; N, 9.06;

Found: C, 66.98; H, 5.67; Cl, 8.06; N, 9.04%.

$[\alpha]_D^{20} = +27.6^\circ$ (C=1.05; CHCl₃).

Example 89

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-methoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 59 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 283-284°C.

Analysis for C₂₂H₂₀ClN₃O₃:

Calculated: C, 64.47; H, 4.92; Cl, 8.65; N, 10.25;

Found: C, 64.49; H, 4.92; Cl, 8.33; N, 10.02%.

$[\alpha]_D^{20} = +61.3^\circ$ (C=1.00; CHCl₃).

Example 90

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3-chloro-4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from isopropylamine and intermediate 59 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 302-304°C.

Analysis for C₂₄H₂₄ClN₃O₃:

Calculated: C, 65.83; H, 5.52; N, 9.60;

Found: C, 65.83; H, 5.57; N, 9.73%.

$[\alpha]_D^{20} = +39.8^\circ$ (C=0.95; CHCl₃).

Example 91

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 61 gave, after recrystallisation from dichloromethane/methanol, the title compound as white crystals m.p.: 288-291°C.

Analysis for C₂₃H₂₁N₃O₃:

Calculated: C, 71.30; H, 5.46; N, 10.85;

Found: C, 71.27; H, 5.49; N, 10.96%.

$[\alpha]_D^{20} = +65.6^\circ$ (C=0.4; CHCl₃).

Example 92

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methylcyclopropylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylcyclopropylamine and intermediate 61 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 242-244°C.

Analysis for $C_{26}H_{25}N_3O_3$:
 Calculated: C, 73.05; H, 5.89; N, 9.83;
 Found: C, 72.90; H, 5.93; N, 9.98%.
 $[\alpha]_D^{20} = +55.4^\circ$ (C=0.99; $CHCl_3$).

5

Example 93

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-indanyl)-2-methylpyrazino(2',1':6,1]pyrido[3,4-b]indole-1,4-dione

10 The same two step procedure but starting from methylamine and intermediate 63 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: $262^\circ C$.

Analysis for $C_{24}H_{23}N_3O_2$:
 Calculated: C, 74.78; H, 6.01; N, 10.90;
 Found: C, 74.65; H, 5.90; N, 10.67%.
 $[\alpha]_D^{20} = +68.6^\circ$ (C=0.98; $CHCl_3$).

15

Example 94

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-indanyl)-2-cyclopropylmethylpyrazino(2',1':6,1]pyrido[3,4-b]indole-1,4-dione

20

The same two step procedure but starting from cyclopropylmethylamine and intermediate 63 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: $176^\circ C$.

Analysis for $C_{27}H_{27}N_3O_2$ ($0.25H_2O$):
 Calculated: C, 75.41; H, 6.45; N, 9.77;
 Found: C, 75.25; H, 6.51; N, 9.75%.
 $[\alpha]_D^{20} = +57.9^\circ$ (C=1.00; $CHCl_3$).

25

Example 95

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino(2',1':6,1]pyrido[3,4-b]indole-1,4-dione

30

To a stirred suspension of Intermediate 73 (12.5g) in MeOH (400ml) was added at room temperature a solution of methylamine (33% in EtOH) (13.7ml) and the resulting mixture was heated at $50^\circ C$ under N_2 for 14 hours. The solvent was removed under reduced pressure and the residue was dissolved in CH_2Cl_2 (1l). After washing with water (3 x 500ml), drying over Na_2SO_4 and evaporating to dryness, the white solid obtained was recrystallised from 2-propanol to give the title compound as white needles (7.5g).
 mp: $298-300^\circ C$.

35

$[\alpha]_D^{20} = +71.3^\circ$ (c = 0.55, $CHCl_3$).
 Elemental analysis ($C_{22}H_{19}N_3O_4$) calculated: C, 67.86; H, 4.92; N, 10.79; found: C, 67.79; H, 4.95; N, 10.61%.

40

Example 96

Cis-2,3,6,7,12,12a-hexahydro-2,10-dimethyl-6-(3,4-methylenedioxyphenyl)-pyrazino(2',1':6,1]pyrido[3,4-b]indole-1,4-dione

45

The same two step procedure as used to prepare Example 1, but starting from methylamine and the cis isomer of Intermediate 74, gave after recrystallisation from ethanol, the title compound as white crystals m.p.: $275^\circ C$.

Analysis for $C_{23}H_{21}N_3O_4$ ($0.4H_2O$):
 Calculated: C, 67.27; H, 5.35; N, 10.23;
 Found: C, 67.36; H, 5.21; N, 10.31%.

50

55

Example 97

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-(3,4-dimethoxybenzyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

5

The same two step procedure as used to prepare Example 78, but starting from veratrylamine and intermediate 54 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 224-226°C.

Analysis for $C_{30}H_{27}N_3O_6$:

Calculated: C, 68.56; H, 5.18; N, 8.00;

10 Found: C, 68.80; H, 5.11; N, 8.06%.

$[\alpha]_D^{20} = +43.9^\circ$ (C = 1.02; $CHCl_3$).

Example 98

15 Cis-2,3,6,7,12,12a-hexahydro-6-(4-aminophenyl)-2-butylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

To a solution of Example 75 (1.5 g) in methanol (100 mL) was added $SnCl_2 \cdot H_2O$ (3.06) and the resulting mixture was heated at reflux for 8 hours. The mixture was cooled to ambient temperature, poured into ice and was adjusted to pH5 with 1N NaOH. The methanol was evaporated off and the residue was basified to pH11 with 1N NaOH and
20 extracted with EtOAc (2 x 150 mL). After drying over Na_2SO_4 and evaporation of EtOAc, the resulting yellow powder was purified by radial chromatography eluting with CH_2Cl_2 to give the title compound as a white powder (550 mg) m.p.: 192°C.

Analysis for $C_{24}H_{26}N_4O_2$ (1.3 H_2O):

Calculated: C, 67.68; H, 6.77; N, 13.15;

25 Found: C, 67.74; H, 6.68; N, 13.02%.

Example 99

30 Cis-2,3,6,7,12,12a-hexahydro-6-(4-acetamidophenyl)-2-butylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

To a solution of Example 98 (0.2 g) in THF (15 mL) was added triethylamine (76 μ L) and acetyl chloride (39 μ L) and the resulting solution was stirred at room temperature for 2 hours. After evaporation of THF, the resulting residue was taken up in CH_2Cl_2 (100 mL), washed with water (2 x 50 mL) and dried over Na_2SO_4 . After evaporation of CH_2Cl_2 , the resulting solid was recrystallised from MeOH/ H_2O to give the title compound as a cream powder (120 mg) m.p.:
35 246°C.

Analysis for $C_{26}H_{28}N_4O_3$:

Calculated: C, 70.25; H, 6.35; N, 12.60;

Found: C, 69.85; H, 6.38; N, 12.56%.

40 Example 100

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methylsulfonamidophenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

45 To a solution of Example 98 (0.2 g) in THF (5 mL) was added triethylamine (228 μ L) and methanesulfonyl chloride (126 μ L) and the solution was heated at reflux for 6 hours. After evaporation of THF, the residue was taken up in CH_2Cl_2 , washed with water and dried over Na_2SO_4 . After evaporation of CH_2Cl_2 , the residue was purified by radial chromatography eluting with CH_2Cl_2 /MeOH (95/5) to give the title compound as a brown powder (30 mg) m.p.: 188°C.

Analysis for $C_{25}H_{28}N_4O_4S$ (0.75 H_2O):

50 Calculated: C, 60.77; H, 6.02; N, 11.34;

Found: C, 60.61; H, 6.02; N, 10.82%.

Example 101 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

55

The same two step procedure but starting from ammonia and intermediate 54 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 285-290°C.

Analysis for $C_{21}H_{17}N_3O_4$:

Calculated : C, 67.19 ; H, 4.56 ; N, 11.19 ;
 Found : C, 67.30 ; H, 4.66 ; N, 11.11 %.
 $[\alpha]^{20}_D = + 88^\circ$ (c = 0.48 ; pyridine).

5 Example 102

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-2-(2-propynyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

10 The same two step procedure but starting from propargylamine and intermediate 54 gave, after recrystallisation from acetone, the title compound as white crystals m.p.: 271°C.

Analysis for $C_{24}H_{19}N_3O_4$:

Calculated : C, 69.72 ; H, 4.63 ; N, 10.16 ;

Found : C, 69.95 ; H, 4.66 ; N, 10.06 %.

15 $[\alpha]^{20}_D = + 51.7^\circ$ (c = 0.49 ; $CHCl_3$).

Example 103

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-(3,4-methylenedioxybenzyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

20 The same two step procedure but starting from piperonylamine and intermediate 54 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 204-206°C.

Analysis for $C_{29}H_{23}N_3O_6$:

25 Calculated : C, 68.36 ; H, 4.55 ; N, 8.25 ;

Found : C, 68.25 ; H, 4.49 ; N, 8.41.

$[\alpha]^{20}_D = + 43^\circ$ (c = 1.01 ; $CHCl_3$).

Example 104

30 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-(3,4-dimethoxyphenethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

35 The same two step procedure but starting from 3,4-dimethoxyphenethylamine and intermediate 54 gave, after recrystallisation from dichloromethane/ether, the title compound as white crystals m.p.: 265-266°C.

Analysis for $C_{31}H_{29}N_3O_6$:

Calculated : C, 69.00 ; H, 5.42 ; N, 7.79 ;

Found : C, 68.68 ; H, 5.35 ; N, 7.78 %.

$[\alpha]^{20}_D = + 38.3^\circ$ (c = 1.12 ; $CHCl_3$).

40

Example 105

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-furfuryl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

45 The same two step procedure but starting from furfurylamine and intermediate 54 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 219°C.

Analysis for $C_{26}H_{21}N_3O_5$:

Calculated : C, 68.56 ; H, 4.65 ; N, 9.23 ;

50 Found : C, 68.16 ; H, 4.63 ; N, 9.15 %.

$[\alpha]^{20}_D = + 58.1^\circ$ (c = 1.2 ; $CHCl_3$)

Example 106

55 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-2-(2-thienylmethyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from 2-thiophenemethylamine and intermediate 54 gave, after recryst-

tallisation from methanol/water, the title compound as white crystals m.p.: 155-157°C.

Analysis for $C_{26}H_{21}N_3O_4S$:

Calculated : C, 66.23 ; H, 4.49 ; N, 8.91 ; S, 6.8 ;

Found : C, 66.13 ; H, 4.54 ; N, 9.12 ; S, 6.78 %.

$[\alpha]^{20}_D = +70.4^\circ$ (c = 1.03 ; $CHCl_3$).

Example 107

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(4-methoxyphenyl)-2-methyl-pyrazino [2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and intermediate 57 gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 285-288°C.

Analysis for $C_{22}H_{21}N_3O_3$:

Calculated : C, 70.38 ; H, 5.64 ; N, 11.19 ;

Found : C, 70.31 ; H, 5.69 ; N, 11.29 %.

$[\alpha]^{20}_D = +59^\circ$ (c = 1.19 ; $CHCl_3$).

Example 108

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-ethyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from ethylamine and intermediate 57 gave, after recrystallisation from methanol, the title compound as white crystals m.p. : 277°C.

Analysis for $C_{23}H_{23}N_3O_3$:

Calculated : C, 70.93 ; H, 5.95 ; N, 10.79 ;

Found : C, 70.90 ; H, 5.96 ; N, 10.54 %.

$[\alpha]^{20}_D = +52^\circ$ (c = 1.28 ; $CHCl_3$).

Example 109

(6R,12aR)-2,3,6,7,12,12a-hexahydro-6-(7-(4-methyl-3,4-dihydro-2H-benzof[1,4]oxazinyl))-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from intermediate 75 and methylamine gave, after recrystallisation from ethanol, the title compound as white crystals m.p.: 285-288°C.

Analysis for $C_{24}H_{24}N_4O_3$ (0.5 H_2O) :

Calculated : C, 67.75 ; H, 5.92 ; N, 13.17 ;

Found : C, 68.02 ; H, 6.00 ; N, 13.18 %.

$[\alpha]^{20}_D = +71.7^\circ$ (c = 1, pyridine).

Example 110

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-(N-benzylindolinyl))-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from intermediate 77 and methylamine gave, after recrystallisation from dichloromethane/methanol, the title compound as white crystals m.p.: 223-225°C.

Analysis for $C_{30}H_{28}N_4O_2$:

Calculated : C, 75.61 ; H, 5.92 ; N, 11.76 ;

Found : C, 75.2 ; H, 5.78 ; N, 11.67 %.

$[\alpha]^{20}_D = +20.4^\circ$ (c = 0.5, $CHCl_3$).

Example 111

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-indolinyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

A solution of Example 110 (1.05 g, 2.2 mmol) in methanol (100 mL) was hydrogenated in the presence of 10 % Pd-C (100 mg) for 48 hours at room temperature. After removal of the catalyst, the solvent was evaporated in vacuo

to leave a residue which was purified by flash chromatography eluting with dichloromethane/methanol : 96/4. The solid obtained was recrystallised from dichloromethane/methanol to give the title compound (300 mg) as white crystals m.p.: 240°C.

Analysis for $C_{23}H_{22}N_4O_2$ (0.5 H_2O) :

5 Calculated : C, 69.86 ; H, 5.86 ; N, 14.17 ;

Found : C, 70.13 ; H, 5.77 ; N, 14.06 %.

$[\alpha]^{20}_D = +55.9^\circ$ (c = 1.18 ; pyridine).

Example 112

10 Cis-2,3,6,7,12,12a-hexahydro-6-(4-ethylphenyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from methylamine and the cis isomer of intermediate 42 gave, after recrystallisation from methanol, the title compound as white crystals m.p. : 254°C.

15 Analysis for $C_{23}H_{23}N_3O_2$ (0.25 H_2O) :

Calculated : C, 73.09 ; H, 6.27 ; N, 11.12 ;

Found : C, 73.03 ; H, 6.18 ; N, 11.36 %.

Example 113

20 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(4-carbomethoxyphenyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two step procedure but starting from intermediate 78 (cis isomer) and methylamine gave, after recrystallisation from methanol, the title compound as white crystals m.p.: 308-312°C.

25 Analysis for $C_{23}H_{21}N_3O_4$:

Calculated : C, 68.47 ; H, 5.25 ; N, 10.42 ;

Found : C, 68.76 ; H, 5.18 ; N, 10.35 %.

$[\alpha]^{20}_D = +97.7^\circ$ (c = 1, pyridine).

Example 114

30 (5aR,12R,14aR)-1,2,3,5a,6,11,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1",2":4',5']pyrazino[2',1':6,1]pyrido[3,4-b]indole-5-1,4-dione

A solution of intermediate 80 (0.7 g, 1.2 mmol) in a mixture of methanol/THF (80/40 mL) was hydrogenated in the presence of 10 % Pd-C (75 mg) for 48 hours at 40°C. After removal of the catalyst, the solvent was evaporated in vacuo to leave a residue, which was purified by flash chromatography eluting with dichloromethane/methanol : 98/2. The white solid obtained was recrystallised from methanol to give the title compound (180 mg) as white crystals m.p.: 284-287°C.

40 Analysis for $C_{24}H_{21}N_3O_4$:

Calculated : C, 69.39 ; H, 5.10 ; N, 10.11 ;

Found : C, 69.47 ; H, 5.11 ; N, 9.97 %.

$[\alpha]^{20}_D = +21.7^\circ$ (c = 0.64, $CHCl_3$).

Example 115

45 (5aR,12R,14aS)-1,2,3,5,6,11,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1",2":4',5']pyrazino[2',1':6,1]pyrido[3,4-b]indole-5-1,4-dione

50 A solution of intermediate 81 (0.8 g, 1.37 mmol) in methanol (40 mL) was hydrogenated in the presence of 10 % Pd-C (100 mg) for 5 h at 45°C. After removal of the catalyst the solvent was evaporated in vacuo to leave a residue, which was purified by flash chromatography eluting with dichloromethane/methanol : 98/2. The solid obtained was recrystallised from methanol to give the title compound (300 mg) as white crystals m.p. : 302-304°C.

55 Analysis for $C_{24}H_{21}N_3O_4$:

Calculated : C, 69.39 ; H, 5.10 ; N, 10.11 ;

Found : C, 69.35 ; H, 5.11 ; N, 10.10 %.

$[\alpha]^{20}_D = +106.8^\circ$ (c = 1.08, $CHCl_3$).

Example 116

(3R, 6R, 12aR)-2,3,6,7,12,12a-hexahydro-2,3-dimethyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

To a stirred solution of intermediate 82 (0.15 g, 0.34 mmol) in THF (15 mL) was added at room temperature a solution of methylamine (33 % in EtOH) (0.32 mL) and the resulting solution was heated at reflux under N₂ for 24 hours. The solvent was removed under reduced pressure and the residue was dissolved in CH₂Cl₂ (25 mL). After washing with water (2 x 20 mL), drying over Na₂SO₄ and evaporating to dryness, the crude product was purified by flash chromatography eluting with dichloromethane/methanol : 99/1. The white solid obtained was recrystallised from methanol to give the title compound as white crystals (80 mg) m.p. : 219-220°C.

Analysis for C₂₃H₂₁N₃O₄ :

Calculated : C, 68.47 ; H, 5.25 ; N, 10.42 ;

Found : C, 68.39; H, 5.21; N, 10.42%.

[α]_D²⁰ = + 89.6° (c = 1 ; CHCl₃).

Example 117

(3S, 6R, 12aR)-2,3,6,7,12,12a-hexahydro-2,3-dimethyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

To a stirred solution of intermediate 83 (0.3 g, 0.68 mmol) in THF (30 mL) was added at room temperature a solution of methylamine (33 % in EtOH) (0.68 mL) and the resulting solution was treated at reflux under N₂ for 6 days. The solvent was removed under reduced pressure and the residue was dissolved in CH₂Cl₂ (50 mL). After washing with water (2,25 mL), drying over Na₂SO₄ and evaporating to dryness, the crude product was purified by flash chromatography eluting with dichloromethane/methanol : 99/1. The oily residue obtained was crystallised from methanol to give the title compound as white crystals (40 mg) m.p. : 307-309°C.

Analysis for C₂₃H₂₁N₃O₄ :

Calculated : C, 68.47 ; H, 5.25 ; N, 10.42 ;

Found : C, 68.35; H, 5.33; N, 10.42%.

[α]_D²⁰ = + 65.2° (c = 1.15 ; CHCl₃).

Example 118

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-dihydroxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

A solution of intermediate 86 (0.75 g ; 1.34 mmol) in a mixture of ethanol/THF (70/30 mL) was hydrogenated in the presence of 10 % Pd-C (75 mg) for 24 h at room temperature. After removal of the catalyst, the solvent was evaporated in vacuo to leave a white solid which was recrystallised from methanol to give the title compound (0.35 g) as white crystals m.p. : 224-226°C.

Analysis for C₂₁H₁₉N₃O₄ :

Calculated : C, 66.83 ; H, 5.07 ; N, 11.13 ;

Found : C, 66.58 ; H, 5.01 ; N, 11.04 %.

[α]_D²⁰ = + 58.4° (c = 1.04 ; pyridine).

Example 119

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(5-(2-methylisindolinyl))pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione

The same two steps procedure but starting from intermediate 87 and methylamine gave a crude oil which was purified by flash chromatography eluting with dichloromethane/methanol/triethylamine : 92/8/0.1 %. The solid obtained was recrystallized from isopropanol/propyl ether/water to give the title compound (20 mg) as off-white crystals m.p.: 236°C.

Analysis for C₂₄H₂₄N₄O₂ (2.68 H₂O)

Calculated : C, 64.23 ; H, 6.59 ; N, 12.48 ;

Found: C, 64.21 ; H, 6.43 ; N, 12.02 %.

EP 0 740 668 B1

$[\alpha]^{20}_D = + 61.1^\circ$ (c = 0.5 ; CH₃OH).

Example 120

Compounds of formula (I) have been included in pharmacy formulations and details of such formulations are given below.

TABLETS FOR ORAL ADMINISTRATION

A. Direct Compression

1.	mg/tablet
Active ingredient	50.0
Crospovidone USNF	8.0
Magnesium Stearate Ph Eur	1.0
Anhydrous Lactose	141.0

The active ingredient was sieved and blended with the excipients. The resultant mix was compressed into tablets.

2.	mg/tablet
Active ingredient	50.0
Colloidal Silicon Dioxide	0.5
Crospovidone	8.0
Sodium Lauryl Sulphate	1.0
Magnesium Stearate Ph Eur	1.0
Microcrystalline Cellulose USNF	139.5

The active ingredient was sieved and blended with the excipients. The resultant mix was compressed into tablets.

B. WET GRANULATION

1.	mg/tablet
Active ingredient	50.0
Polyvinyl pyrrolidone	150.0
Polyethylene glycol	50.0
Polysorbate 80	10.0
Magnesium Stearate Ph Eur	2.5
Croscarmellose Sodium	25.0
Colloidal Silicon Dioxide	2.5
Microcrystalline Cellulose USNF	210.0

The polyvinyl pyrrolidone, polyethylene glycol and polysorbate 80 were dissolved in water. The resultant solution was used to granulate the active ingredient. After drying the granules were screened, then extruded at elevated temperatures and pressures. The extrudate was milled and/or screened then was blended with the microcrystalline cellulose, croscarmellose sodium, colloidal silicon dioxide and magnesium stearate. The resultant mix was compressed into tablets.

2.	mg/tablet mg/tablet
Active ingredient	50.0
Polysorbate 80	3.0
Lactose Ph Eur	178.0
Starch BP	45.0

EP 0 740 668 B1

(continued)

2.	mg/tablet mg/tablet
Pregelatinised Maize Starch BP	22.5
Magnesium Stearate BP	1.5

The active ingredient was sieved and blended with the lactose, starch and pregelatinised maize starch. The polysorbate 80 was dissolved in purified water. Suitable volumes of the polysorbate 80 solution were added and the powders were granulated. After drying, the granules were screened and blended with the magnesium stearate. The granules were then compressed into tablets.

Tablets of other strengths may be prepared by altering the ratio of active ingredient to the other excipients.

FILM COATED TABLETS

The aforementioned tablet formulations were film coated.

Coating Suspension	% w/w
Opadry white†	13.2
Purified water Ph Eur	to 100.0*

* The water did not appear in the final product. The maximum theoretical weight of solids applied during coating was 20mg/tablet.

†Opadry white is a proprietary material obtainable from Colorcon Limited, UK which contains hydroxypropyl methylcellulose, titanium dioxide and triacetin.

The tablets were film coated using the coating suspension in conventional film coating equipment.

CAPSULES

1.	mg/capsule
Active ingredient	50.0
Lactose	148.5
Polyvinyl pyrrolidone	100.0
Magnesium Stearate	1.5

The active ingredient was sieved and blended with the excipients. The mix was filled into size No. 1 hard gelatin capsules using suitable equipment.

2.	mg/capsule
Active ingredient	50.0
Microcrystalline Cellulose	233.5
Sodium Lauryl Sulphate	3.0
Crospovidone	12.0
Magnesium Stearate	1.5

The active ingredient was sieved and blended with the excipients. The mix was filled into size No. 1 hard gelatin capsules using suitable equipment.

Other doses may be prepared by altering the ratio of active ingredient to excipient, the fill weight and if necessary changing the capsule size.

3.	mg/capsule
Active ingredient	50.0
Labrafil M1944CS	to 1.0 ml

The active ingredient was sieved and blended with the Labrafil. The suspension was filled into soft gelatin capsules using appropriate equipment.

Example 121

Inhibitory effect on cGMP-PDE

cGMP-PDE activity of compounds of the present invention was measured using a one-step assay adapted from Wells et al. (Wells, J. N., Baird, C. E., Wu, Y. J. and Hardman, J. G., Biochim. Biophys. Acta 384, 430 (1975)). The reaction medium contained 50mM Tris-HCl, pH 7.5, 5mM Mg-acetate, 250µg/ml 5'-Nucleotidase, 1mM EGTA and 0.15µM 8-[H³]-cGMP. The enzyme used was a human recombinant PDE V (ICOS, Seattle USA).

Compounds of the invention were dissolved in DMSO finally present at 2% in the assay. The incubation time was 30 minutes during which the total substrate conversion did not exceed 30%.

The IC₅₀ values for the compounds examined were determined from concentration-response curves using typically concentrations ranging from 10nM to 10µM. Tests against other PDE enzymes using standard methodology also showed that compounds of the invention are highly selective for the cGMP specific PDE enzyme.

-cGMP level measurements

Rat aortic smooth muscle cells (RSMC) prepared according to Chamley et al. in Cell Tissue Res. 177, 503 - 522 (1977) were used between the 10th and 25th passage at confluence in 24-well culture dishes. Culture media was aspirated and replaced with PBS (0.5ml) containing the compound tested at the appropriate concentration. After 30 minutes at 37°C, particulate guanylate cyclase was stimulated by addition of ANF (100nM) for 10 minutes. At the end of incubation, the medium was withdrawn and two extractions were performed by addition of 65% ethanol (0.25ml). The two ethanolic extracts were pooled and evaporated until dryness, using a Speed-vac system. c-GMP was measured after acetylation by scintillation proximity immunoassay (AMERSHAM).

The compounds according to the present invention were typically found to exhibit an IC₅₀ value of less than 500nM, and an EC₅₀ value of less than 5. In vitro test data for representative compounds of the invention is given in following Table 1:

Table 1

Example No.	IC ₅₀ nM	EC ₅₀ µM
12	10	0.15
36	<10	0.5
52	20	0.8
63	30	0.35
79	<10	0.15
82	20	0.5
84	10	0.4
89	10	<0.1
95	2	0.2
101	10	0.3
115	<10	0.4

Example 122

-Antihypertensive activity in rats

The hypotensive effects of compounds according to the invention as identified in table 2 were studied in conscious spontaneously hypertensive rats (SHR). The compounds were administered orally at a dose of 5mg/kg in a mixture of 5% DMF and 95% olive oil. Blood pressure was measured from a catheter inserted in the carotid artery and recorded for 5 hours after administration. The results are expressed as Area Under the Curve (AUC from 0 to 5 hours, mmHg.

hour) of the fall in blood pressure over time.

In Vivo Results

5

10

15

20


Example No.	AUC PO (mmHg.h)
95	135
101	136
36	99
63	95
79	171
82	111
84	77
89	117
95	135
101	136

Claims

25

1. A compound of formula (I)

30



35

and salts and solvates thereof, in which:

40


R⁰ represents hydrogen, halogen or C₁₋₆ alkyl;

R¹ represents hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆ alkynyl, haloC₁₋₆alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₃alkyl, arylC₁₋₃alkyl or heteroarylC₁₋₃alkyl, where aryl means phenyl or phenyl substituted by one or more (e.g. 1, 2 or 3) substituents selected from halogen, C₁₋₆alkyl, C₁₋₆alkoxy and methylenedioxy, and heteroaryl means thienyl, furyl or pyridyl each optionally substituted by one or more (e.g. 1, 2 or 3) substituents selected from halogen, C₁₋₆ alkyl and C₁₋₆alkoxy;

45

R² represents a monocyclic aromatic ring selected from benzene, optionally substituted by one or more (e.g. 1, 2 or 3) atoms or groups comprising halogen, hydroxy, C₁₋₆alkyl, C₁₋₆alkoxy, -CO₂R^b, haloC₁₋₆alkyl, haloC₁₋₆alkoxy, cyano, nitro and NR^aR^b, or R² represents an optionally substituted thiophene, furan, pyridine, or a bicyclic ring

50



55


attached to the rest of the molecule via one of the benzene ring carbon atoms and wherein the fused ring A is a 5- or 6-membered ring which may be saturated or partially or fully unsaturated and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulphur and nitrogen; where optional

substitution means one or more (e.g. 1, 2 or 3) atoms or groups comprising halogen, C₁₋₆alkyl, C₁₋₆alkoxy and arylC₁₋₃alkyl as defined above;

R³ represents hydrogen or C₁₋₃ alkyl, or R¹ and R³ together represent a 3- or 4-membered alkyl or alkenyl chain; and

R^a and R^b are each hydrogen or C₁₋₆alkyl, or R^a may also represent C₂₋₇alkanoyl or C₁₋₆alkylsulphonyl.

2. A compound of formula (Ia)




and salts and solvates thereof, in which:

R⁰ represents hydrogen, halogen or C₁₋₆ alkyl;

R¹ represents hydrogen, C₁₋₆alkyl, haloC₁₋₆alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₃alkyl, arylC₁₋₃alkyl or heteroarylC₁₋₃alkyl, where aryl means phenyl or phenyl substituted by one or more (e.g. 1, 2 or 3) substituents selected from halogen, C₁₋₆alkyl, C₁₋₆alkoxy and methylenedioxy, and heteroaryl means thienyl, furyl or pyridyl each optionally substituted by one or more (e.g. 1, 2 or 3) substituents selected from halogen, C₁₋₆ alkyl and C₁₋₆alkoxy;

R² represents a monocyclic aromatic ring selected from benzene, optionally substituted by one or more (e.g. 1, 2 or 3) atoms or groups comprising halogen, hydroxy, C₁₋₆alkyl, C₁₋₆alkoxy, -CO₂R^b, haloC₁₋₆alkyl, haloC₁₋₆alkoxy, cyano, nitro and NR^aR^b, or R² represents an optionally substituted thiophene, furan, pyridine, or a bicyclic ring



attached to the rest of the molecule via one of the benzene ring carbon atoms and wherein the fused ring A is a 5- or 6-membered ring which may be saturated or partially or fully unsaturated and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulphur and nitrogen, where optional substitution means one or more (e.g. 1, 2 or 3) atoms or groups halogen, C₁₋₆alkyl, C₁₋₆alkoxy and arylC₁₋₃alkyl as defined above; and

R^a and R^b are each hydrogen or C₁₋₆alkyl, or R^a may also represent C₂₋₇alkanoyl or C₁₋₆alkylsulphonyl.


3. A compound according to Claim 1 or 2, wherein R⁰ represents hydrogen.

4. A compound according to any of Claims 1 to 3, wherein R¹ represents hydrogen, C₁₋₄alkyl, haloC₁₋₄alkyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkylmethyl, pyridyl C₁₋₃alkyl, furylC₁₋₃alkyl or optionally substituted benzyl.

5. A compound according to Claim 1, wherein R¹ and R³ together represent a 3-membered alkyl chain.


6. A compound according to Claim 1, wherein R³ represents hydrogen.

7. A compound according to any of Claims 1 to 6, wherein R² represents an optionally substituted benzene, thiophene, furan, pyridine or naphthalene ring or an optionally substituted bicyclic ring



where n is 1 or 2 and X and Y are each CH₂ or O.

8. A cis isomer of formula (I) represented by formula (Ib)



and mixtures thereof with its cis optical enantiomer, including racemic mixtures, and salts and solvates of these compounds in which R⁰ is hydrogen or halogen and R¹, R² and R³ are as defined in any preceding claim.

9. Cis-2,3,6,7,12,12a-hexahydro-2-(4-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thienyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methylphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-methoxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-methoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(5aR,12R,14aS)-1,2,3,5,6,1,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1',2'':4',5']pyrazino[2',1':6,1]pyrido[3,4-b]indole-5-1,4-dione;

and physiologically acceptable salts and solvates thereof.

10. (6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione; and physiologically acceptable salts and solvates thereof.

11. A compound according to any of Claims 1 to 10, for use in the treatment of stable, unstable and variant angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, congestive heart failure, renal failure, atherosclerosis, conditions of reduced blood vessel patency, peripheral vascular disease, vascular disorders inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility.


12. Use of a compound according to any of Claims 1 to 10, for the manufacture of a medicament for the treatment of stable, unstable and variant angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, congestive heart failure, renal failure, atherosclerosis, conditions of reduced blood vessel patency, peripheral vascular disease, vascular disorders inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility.

13. A pharmaceutical composition comprising a compound of the according to any of Claims 1 to 10, together with a pharmaceutically acceptable diluent or carrier therefor.

14. A process of preparing a pharmaceutical composition comprising a compound according to any of Claims 1 to 10, which process comprises mixing said compound together with a pharmaceutically acceptable diluent or carrier therefor.


5 15. A process of preparing a compound of formula (I), which process comprises:

a process (A) for preparing a compound of formula (I), wherein R^3 represents hydrogen which process (A) comprises treating a compound of formula (II)




in which Alk represents C_{1-6} alkyl and Hal is a halogen atom, with a primary amine R^1NH_2 ; or

20 a process (B) for preparing a compound of formula (I), wherein R^1 and R^3 together represent a 3- or 4-membered alkyl or alkenyl chain, which process (B) comprises cyclisation of a compound of formula (VIII)



wherein Alk represents C_{1-6} alkyl and R^1 and R^3 together represent a 3- or 4-membered chain both as defined above; or

35 a process (C) for preparing a compound of formula (I) wherein R^3 represents C_{1-3} alkyl, which process (C) comprises cyclisation of a compound of formula (X)




wherein Alk represents C_{1-6} alkyl and R^5 represents C_{2-5} alkyl, substituted at C_1 by a halogen atom; or process (A), (B) or (C) as hereinbefore described followed by


- 50
- i) an interconversion step; and/or either
 - ii) salt formation; or
 - iii) solvate formation.

55 16. Compounds of formulae (II), (III), (V), (VI), (VII), (VIII) and (X)

5




10




15

20




25

30




35

40




45

50




55



10 where R^0 and R^2 are hereinbefore defined as in Claim 1, R^1 and R^3 together represent a 3- or 4-membered alkyl or alkenyl chain, R^5 represents C_{2-6} alkyl substituted at C_1 by a halogen atom, Alk represents C_{1-6} alkyl and Hal is a halogen atom, with the exception of compounds (III), (V), (VI) and (VII) wherein R^0 is hydrogen, R^2 is phenyl and Alk is methyl.

15 **Patentansprüche**

1. Eine Verbindung der Formel (I)




und Salze und Solvate derselben, in der:

30 R^0 Wasserstoff, Halogen oder C_{1-6} -Alkyl darstellt;

35 R^1 Wasserstoff, C_{1-6} -Alkyl, C_{2-6} -Alkenyl, C_{2-6} -Alkinyl, Halo- C_{1-6} -Alkyl, C_{3-8} -Cycloalkyl, C_{3-8} -Cycloalkyl- C_{1-3} -Alkyl, Aryl- C_{1-3} -Alkyl und Heteroaryl- C_{1-3} -Alkyl darstellt, wobei Aryl Phenyl bedeutet oder Phenyl, substituiert mit einem oder mehreren (z.B. 1, 2 oder 3) Substituenten, die ausgewählt sind aus Halogen, C_{1-6} -Alkyl, C_{1-6} -Alkoxy und Methylendioxy, und Heteroaryl Thienyl, Furyl oder Pyridyl bedeutet, jedes fakultativ substituiert mit einem oder mehreren (z.B. 1, 2 oder 3) Substituenten, die ausgewählt sind aus Halogen, C_{1-6} -Alkyl und C_{1-6} -Alkoxy;

40 R^2 einen monocyclischen aromatischen Ring darstellt, ausgewählt aus Benzol, fakultativ substituiert mit einem oder mehreren (z.B. 1, 2 oder 3) Atomen oder Gruppen, die Halogen, Hydroxy, C_{1-6} -Alkyl, C_{1-6} -Alkoxy, $-CO_2R^b$, Halo- C_{1-6} -Alkyl, Halo- C_{1-6} -Alkoxy, Cyano, Nitro und NR^aR^b umfassen, oder R^2 einen fakultativ substituierten Thiophen-, Furan-, Pyridin-Ring darstellt oder einen bicyclischen Ring




50 der an den Rest des Moleküls über eines der Benzolring-Kohlenstoffatome gebunden ist, und wobei der anellierte Ring A ein 5- oder 6-gliedriger Ring ist, der gesättigt oder teilweise oder vollständig ungesättigt sein kann und Kohlenstoffatome und fakultativ ein oder zwei Heteroatome umfaßt, die ausgewählt sind aus Sauerstoff, Schwefel und Stickstoff; wobei fakultative Substitution ein oder mehrere (z.B. 1, 2 oder 3) Atome oder Gruppen bedeutet, die Halogen, C_{1-6} -Alkyl, C_{1-6} -Alkoxy und Aryl- C_{1-3} -Alkyl umfassen, wie oben definiert;

55 R^3 Wasserstoff oder C_{1-3} -Alkyl darstellt oder R^1 und R^3 zusammen eine 3- oder 4-gliedrige Alkyl- oder Alkenylkette darstellen; und

R^a und R^b jeweils Wasserstoff oder C_{1-6} -Alkyl sind oder R^a auch C_{2-7} -Alkanoyl oder C_{1-6} -Alkylsulfonyl darstellen kann.

2. Eine Verbindung der Formel (Ia)




und Salze und Solvate derselben, in der:

R^0 Wasserstoff, Halogen oder C_{1-6} -Alkyl darstellt;

R^1 Wasserstoff, C_{1-6} -Alkyl, Halo- C_{1-6} -Alkyl, C_{3-8} -Cycloalkyl, C_{3-8} -Cycloalkyl- C_{1-3} -Alkyl, Aryl- C_{1-3} -Alkyl oder Heteroaryl- C_{1-3} -Alkyl darstellt, wobei Aryl Phenyl bedeutet oder Phenyl, substituiert mit einem oder mehreren (z.B. 1, 2 oder 3) Substituenten, die ausgewählt sind aus Halogen, C_{1-6} -Alkyl, C_{1-6} -Alkoxy und Methylendioxy, und Heteroaryl Thienyl, Furyl oder Pyridyl bedeutet, jedes fakultativ substituiert mit einem oder mehreren (z.B. 1, 2 oder 3) Substituenten, die ausgewählt sind aus Halogen, C_{1-6} -Alkyl und C_{1-6} -Alkoxy;

R^2 einen monocyclischen aromatischen Ring darstellt, ausgewählt aus Benzol, fakultativ substituiert mit einem oder mehreren (z.B. 1, 2 oder 3) Atomen oder Gruppen, die Halogen, Hydroxy, C_{1-6} -Alkyl, C_{1-6} -Alkoxy, $-CO_2R^b$, Halo- C_{1-6} -Alkyl, Halo- C_{1-6} -Alkoxy, Cyano, Nitro und NR^aR^b umfassen, oder R^2 einen fakultativ substituierten Thiophen-, Furan-, Pyridin-Ring darstellt oder einen bicyclischen Ring



der an den Rest des Moleküls über eines der Benzolring-Kohlenstoffatome gebunden ist, und wobei der anellierte Ring A ein 5- oder 6-gliedriger Ring ist, der gesättigt oder teilweise oder vollständig ungesättigt sein kann und Kohlenstoffatome und fakultativ ein oder zwei Heteroatome umfaßt, die ausgewählt sind aus Sauerstoff, Schwefel und Stickstoff, wobei fakultative Substitution ein oder mehrere (z.B. 1, 2 oder 3) Atome oder Gruppen Halogen, C_{1-6} -Alkyl, C_{1-6} -Alkoxy und Aryl- C_{1-3} -Alkyl bedeutet, wie oben definiert; und

R^a und R^b jeweils Wasserstoff oder C_{1-6} -Alkyl sind oder R^a auch C_{2-7} -Alkanoyl oder C_{1-6} -Alkylsulfonyl darstellen kann.


3. Eine Verbindung nach Anspruch 1 oder 2, wobei R^0 Wasserstoff darstellt.

4. Eine Verbindung nach einem der Ansprüche 1 bis 3, wobei R^1 Wasserstoff, C_{1-4} -Alkyl, Halo- C_{1-4} -Alkyl, C_{3-6} -Cycloalkyl, C_{3-6} -Cycloalkylmethyl, Pyridyl- C_{1-3} -Alkyl, Furyl- C_{1-3} -Alkyl oder fakultativ substituiertes Benzyl darstellt.

5. Eine Verbindung nach Anspruch 1, wobei R^1 und R^3 zusammen eine 3-gliedrige Alkylkette darstellen.


6. Eine Verbindung nach Anspruch 1, wobei R^3 Wasserstoff darstellt.

7. Eine Verbindung nach einem der Ansprüche 1 bis 6, wobei R^2 einen fakultativ substituierten Benzol-, Thiophen-, Furan-, Pyridin- oder Naphthalin-Ring darstellt oder einen fakultativ substituierten bicyclischen Ring



worin n 1 oder 2 ist und X und Y jeweils CH₂ oder O sind.

8. Ein cis-Isomer von Formel (I), dargestellt durch Formel (Ib)



und Mischungen desselben mit seinem optischen cis-Enantiomer, einschließlich racemischer Mischungen, und Salze und Solvate dieser Verbindungen, in der R⁰ Wasserstoff oder Halogen ist und R¹, R² und R³ wie in einem vorangehenden Anspruch definiert sind.

9. cis-2,3,6,7,12,12a-Hexahydro-2-(4-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

cis-2,3,6,7,12,12a-Hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

cis-2,3,6,7,12,12a-Hexahydro-6-(5-brom-2-thienyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

cis-2,3,6,7,12,12a-Hexahydro-2-butyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chlor-4-methoxyphenyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

(5aR,12R,14aS)-1,2,3,5,6,11,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1'',2'':4',5']pyrazino[2',1':6,1]pyrido[3,4-b]indol-5-1,4-dion;

und physiologisch annehmbare Salze und Solvate derselben.

10. (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indol-1,4-dion;

und physiologisch annehmbare Salze und Solvate derselben.


11. Eine Verbindung nach einem der Ansprüche 1 bis 10, zur Verwendung bei der Behandlung von stabiler, instabiler und varianter Angina, Bluthochdruck, pulmonarem Bluthochdruck, chronischer obstruktiver Lungenerkrankung, kongestiver Herz insuffizienz, Nierenversagen, Atherosklerose, Zuständen verringerter Blutgefäßdurchgängigkeit, Periphergefäßerkrankung, Gefäßstörungen, entzündlichen Erkrankungen, Schlaganfall, Bronchitis, chronischem Asthma, allergischem Asthma, allergischer Rhinitis, Glaucom oder Erkrankungen, die durch Störungen der Darm-

motilität gekennzeichnet sind.

12. Verwendung einer Verbindung nach einem der Ansprüche 1 bis 10 zur Herstellung eines Medikaments zur Behandlung von stabiler, instabiler und varianter Angina, Bluthochdruck, pulmonarem Bluthochdruck, chronischer obstruktiver Lungenerkrankung, kongestiver Herzinsuffizienz, Nierenversagen, Atherosklerose, Zuständen verringerter Blutgefäßdurchgängigkeit, Periphergefäßerkrankung, Gefäßstörungen, entzündlichen Erkrankungen, Schlaganfall, Bronchitis, chronischem Asthma, allergischem Asthma, allergischer Rhinitis, Glaucom oder Erkrankungen, die durch Störungen der Darmmotilität gekennzeichnet sind.
13. Eine pharmazeutische Zusammensetzung, die eine Verbindung nach einem der Ansprüche 1 bis 10 umfaßt, zusammen mit einem pharmazeutisch annehmbaren Verdünnungsmittel oder Trägerstoff dafür.
14. Ein Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, die eine Verbindung nach einem der Ansprüche 1 bis 10 umfaßt, wobei das Verfahren das Zusammenmischen besagter Verbindung mit einem pharmazeutisch annehmbaren Verdünnungsmittel oder Trägerstoff dafür umfaßt.
15. Ein Verfahren zur Herstellung einer Verbindung von Formel (I), wobei das Verfahren umfaßt:


ein Verfahren (A) zur Herstellung einer Verbindung von Formel (I),

in der R³ Wasserstoff darstellt, wobei das Verfahren (A) das Behandeln einer Verbindung von Formel (II)




in der Alk C₁₋₆-Alkyl darstellt und Hal ein Halogenatom ist, mit einem primären Amin R¹NH₂ umfaßt; oder

ein Verfahren (B) zur Herstellung einer Verbindung von Formel (I), in der R¹ und R³ zusammen eine 3- oder 4-gliedrige Alkyl- oder Alkenylkette darstellen, wobei das Verfahren (B) die Cyclisierung einer Verbindung von Formel (VIII) umfaßt



in der Alk C₁₋₆-Alkyl darstellt und R¹ und R³ zusammen eine 3- oder 4-gliedrige Kette darstellen, beides wie oben definiert; oder

ein Verfahren (C) zur Herstellung einer Verbindung von Formel (I), in der R³ C₁₋₃-Alkyl darstellt, wobei das Verfahren (C) die Cyclisierung einer Verbindung von Formel (X) umfaßt



10 in der Alk C₁₋₆-Alkyl darstellt und R⁵ C₂₋₅-Alkyl darstellt, substituiert am C₁ durch ein Halogenatom; oder




Verfahren (A), (B) oder (C), wie hierin zuvor beschrieben, gefolgt von




15 i) einem Interkonversionsschritt; und/oder entweder

ii) Salzbildung; oder

iii) Solvatbildung.

16. Verbindungen der Formeln (II), (III), (V), (VI), (VII), (VIII) und (X)





20


25

30

in denen R^0 und R^2 hierin zuvor wie in Anspruch 1 definiert sind, R^1 und R^3 zusammen eine 3- oder 4-gliedrige Alkyl- oder Alkenylkette darstellen, R^5 C_{2-5} -Alkyl darstellt, substituiert am C_1 durch ein Halogenatom, Alk C_{1-6} -Alkyl darstellt und Hal ein Halogenatom ist, mit der Ausnahme von Verbindungen (III), (V), (VI) und (VII), bei denen R^0 Wasserstoff ist, R^2 Phenyl ist und Alk Methyl ist.

35 Revendications

1. Composé de formule (I) :




et sels et solvates de celui-ci, dans laquelle :

50 R^0 représente un atome d'hydrogène, d'halogène ou un groupe alkyle en C_{1-6} ;

R^1 représente un atome d'hydrogène, un groupe alkyle en C_{1-6} , alcényle en C_{2-6} , halo-alkyle en C_{1-6} , cycloalkyle en C_{3-8} , cycloalkyl-(en C_{3-8})-alkyle en C_{1-3} , arylalkyle en C_{1-3} , ou hétéroarylalkyle en C_{1-3} , où un groupe aryle signifie un groupe phényle ou phényle substitué par un ou plusieurs (par exemple, 1, 2 ou 3) substituants choisis parmi un atome d'halogène, un groupe alkyle en C_{1-6} , alcoxy en C_{1-6} et méthylènedioxy, et un groupe hétéroaryle signifie un groupe thiényle, furyle ou pyridyle chacun de ceux-ci étant éventuellement substitué par un ou plusieurs (par exemple, 1, 2 ou 3) substituants choisis parmi un atome d'halogène, un groupe alkyle en C_{1-6} et alcoxy en C_{1-6} ;

R^2 représente un noyau aromatique monocyclique choisi parmi le benzène éventuellement substitué par un ou

plusieurs (par exemple, 1, 2 ou 3) atomes ou groupes comprenant un atome d'halogène, un groupe hydroxy, alkyle en C₁₋₆, alcoxy en C₁₋₆, -CO₂R^b, haloalkyle en C₁₋₆, haloalcoxy en C₁₋₆, cyano, nitro et NR^aR^b, ou bien R² représente un groupe thiophène, furanne, pyridine éventuellement substitué, ou noyau bicyclique




un attaché au reste de la molécule par l'intermédiaire d'un des atomes de carbone du cycle benzène et dans lequel le noyau condensé A est un noyau à 5 ou 6 chaînons qui peut être saturé ou partiellement ou complètement insaturé et comprend des atomes de carbone et éventuellement un ou deux hétéroatomes choisis parmi les atomes d'oxygène, de soufre et d'azote; où une substitution éventuelle signifie un ou plusieurs (par exemple, 1, 2 ou 3) atomes ou groupes comprenant un atome d'halogène, un groupe alkyle en C₁₋₆, alcoxy en C₁₋₆ et arylalkyle en C₁₋₃ comme cela est défini plus haut;

R³ représente un atome d'hydrogène ou un groupe alkyle en C₁₋₃, ou bien R¹ et R³ représentent ensemble une chaîne alkyle ou alcényle à 3 ou 4 chaînons; et

R^a et R^b sont chacun un atome d'hydrogène ou un groupe alkyle en C₁₋₆, ou bien R^a peut aussi représenter un groupe alcanoyle en C₂₋₇ ou alkylsulfonyl en C₁₋₆.

2. Composé de formule (Ia) :




et sels et solvates de celui-ci, dans laquelle :

R⁰ représente un atome d'hydrogène, un atome d'halogène ou un groupe alkyle en C₁₋₆;

R¹ représente un atome d'hydrogène, un groupe alkyle en C₁₋₆, halo-alkyle en C₁₋₆, cycloalkyle en C₃₋₈, cycloalkyl-(en C₃₋₈)-alkyle en C₁₋₃, arylalkyle en C₁₋₃, ou hétéroarylalkyle en C₁₋₃, où un groupe aryle signifie un groupe phényle ou phényle substitué par un ou plusieurs (par exemple, 1, 2 ou 3) substituants choisis parmi un atome d'halogène, un groupe alkyle en C₁₋₆, alcoxy en C₁₋₆ et méthylènedioxy, et un groupe hétéroaryle signifie un groupe thiényl, furyl ou pyridyl chacun d'entre eux étant éventuellement substitué par un ou plusieurs (par exemple, 1, 2 ou 3) substituants choisis parmi un atome d'halogène, un groupe alkyle en C₁₋₆ et alcoxy en C₁₋₆;


R² représente un noyau aromatique monocyclique choisi parmi le benzène éventuellement substitué par un ou plusieurs (par exemple, 1, 2 ou 3) atomes ou groupes comprenant un atome d'halogène, un groupe hydroxy, alkyle en C₁₋₆, alcoxy en C₁₋₆, -CO₂R^b, haloalkyle en C₁₋₆, haloalcoxy en C₁₋₆, cyano, nitro et NR^aR^b, ou bien R² représente un groupe thiophène, furanne, pyridine éventuellement substitué, ou noyau bicyclique



un attaché au reste de la molécule par l'intermédiaire d'un des atomes de carbone du cycle benzène et dans lequel le noyau condensé A est un noyau à 5 ou 6 chaînons qui peut être saturé ou partiellement ou complètement insaturé et comprend des atomes de carbone et éventuellement un ou deux hétéroatomes choisis parmi les atomes d'oxygène, de soufre et d'azote; où une substitution éventuelle signifie un ou plusieurs (par


exemple, 1, 2 ou 3) atomes ou groupes comprenant un atome d'halogène, un groupe alkyle en C₁₋₆, alcoxy en C₁₋₆ et arylalkyle en C₁₋₃ comme cela est défini plus haut; et R^a et R^b sont chacun un atome d'hydrogène ou un groupe alkyle en C₁₋₆, ou bien R^a peut aussi représenter un groupe alcanoylé en C₂₋₇ ou alkylsulfonylé en C₁₋₆.

3. Composé suivant les revendications 1 ou 2, dans lequel R⁰ représente un atome d'hydrogène.
4. Composé suivant l'une quelconque des revendications 1 à 3, dans lequel R¹ représente un atome d'hydrogène, un groupe alkyle en C₁₋₄, halo-alkyle en C₁₋₄, cycloalkyle en C₃₋₆, cycloalkyl-(en C₃₋₆)-méthyle, pyridylalkyle en C₁₋₃, furylalkyle en C₁₋₃, ou benzyle éventuellement substitué.
5. Composé suivant la revendication 1, dans lequel R¹ et R³ représentent ensemble une chaîne alkyle à 3 chaînons.
6. Un composé selon la revendication 1, dans lequel R³ représente un atome d'hydrogène.
7. Composé suivant l'une quelconque des revendications 1 à 6, dans lequel R² représente un noyau benzène, thiophène, furanne, pyridine ou naphthalène éventuellement substitué ou un noyau bicyclique éventuellement substitué



dans lequel n est 1 ou 2, et X et Y sont chacun un groupe CH₂ ou O.

8. Isomère cis de formule (I) représenté par la formule (Ib) :



et mélanges de celui-ci avec son énantiomère optique cis, y compris des mélanges racémiques, et sels et solvates de ces composés, dans laquelle R⁰ est un atome d'hydrogène ou d'halogène et R¹, R² et R³ sont tels que définis dans l'une quelconque des revendications précédentes.

9. Cis-2,3,6,7,12,12a-hexahydro-2-(4-pyridylméthyl)-6-(3,4-méthylènedioxyphényl)-pyrazino-[2',1':6,1]-pyrido-[3,4-b]-indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-6-(2,3-dihydrobenzo-[b]-furan-5-yl)-2-méthylpyrazino-[2',1':6,1]-pyrido-[3,4-b]-indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thiényl)-2-méthylpyrazino-[2',1':6,1]-pyrido-[3,4-b]-indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-méthylphényl)-pyrazino-[2',1':6,1]-pyrido-[3,4-b]-indole-1,4-dione;

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-méthylène-dioxyphényl)-pyrazino-[2',1':6,1]-pyrido-[3,4-b]-indole-1,4-dione;

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-méthylène-dioxyphényl)-pyrazino-[2',1':6,1]-pyrido-[3,4-b]-indole-1,4-dione;

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-méthoxyphényl)-pyrazino-[2',1':6,1]-pyrido-[3,4-b]-indole-1,4-dione;

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-méthoxyphényl)-2-méthylpyrazino-[2',1':6,1]-pyrido-

[3,4-b]-indole-1,4-dione;
 (6R, 12aR)-2,3,6,7,12,12a-Hexahydro-2-méthyl-6-(3,4-méthylène-dioxyphényl)-pyrazino-[2',1':6,1]-pyrido-
 [3,4-b]-indole-1,4-dione;
 (6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-méthylènedioxyphényl)-pyrazino-[2',1':6,1]-pyrido-[3,4-b]-indo-
 le-1,4-dione;
 (5aR, 12R, 14aS)-1,2,3,5,6,11,12,14a-Octahydro-12-(3,4- méthylène-dioxyphényl)-pyrrolo-[1'',2'':4',5']-pyra-
 zino
 [2',1':6,1]-pyrido-[3,4-b]-indole-5-1,4-dione;

et sels et solvates de celles-ci acceptables du point de vue physiologique.

10. (6R, 12aR)-2,3,6,7,12,12a-Hexahydro-2-méthyl-6-(3,4-méthylènedioxyphényl)-pyrazino-[2',1':6,1]-pyrido-[3,4-b]-indole-1,4-dione; et sels et solvates de celle-ci acceptables du point de vue physiologique.

11. Composé suivant l'une quelconque des revendications 1 à 10, utile dans le traitement d'angor stable, instable ou type Prinzmetal, d'hypertension, d'hypertension pulmonaire, d'affection pulmonaire obstructive chronique, d'insuffisance cardiaque congestive, d'insuffisance rénale, d'athérosclérose, de situations d'état ouvert réduit de vaisseaux sanguins, d'affection vasculaire périphérique, de troubles vasculaires, d'affections inflammatoires, d'accident vasculaire cérébral constitué, de bronchite, d'asthme chronique, d'asthme allergique, de rhinite allergique, de glaucome ou d'affections caractérisées par des troubles de la motilité intestinale.


12. Utilisation d'un composé suivant l'une quelconque des revendications 1 à 10, pour la fabrication d'un médicament pour le traitement d'angor stable, instable ou type Prinzmetal, d'hypertension, d'hypertension pulmonaire, d'affection pulmonaire obstructive chronique, d'insuffisance cardiaque congestive, d'insuffisance rénale, d'athérosclérose, de situations d'état ouvert réduit de vaisseaux sanguins, d'affection vasculaire périphérique, de troubles vasculaires, d'affections inflammatoires, d'accident vasculaire cérébral constitué, de bronchite, d'asthme chronique, d'asthme allergique, de rhinite allergique, de glaucome ou d'affections caractérisées par des troubles de la motilité intestinale.

13. Composition pharmaceutique comprenant un composé suivant l'une quelconque des revendications 1 à 10, ainsi qu'un diluant ou un support pour celui-ci acceptable du point de vue pharmaceutique.

14. Procédé de préparation d'une composition pharmaceutique comprenant un composé suivant l'une quelconque des revendications 1 à 10, lequel procédé comprend le mélange de ce composé avec un diluant ou un support pour celui-ci acceptable du point de vue pharmaceutique.


15. Procédé de préparation d'un composé de formule (I), lequel procédé comprend :

un procédé (A) pour préparer un composé de formule (I) dans laquelle R³ représente un atome d'hydrogène, lequel procédé (A) comprend le traitement d'un composé de formule (II) :




dans laquelle Alk représente un groupe alkyle en C₁₋₆ et Hal est un atome d'halogène, avec une amine primaire R¹NH₂; ou

un procédé (B) pour préparer un composé de formule (I) dans laquelle R¹ et R³ représentent ensemble une chaîne alkyle ou alcényle à 3 ou 4 chaînons, lequel procédé (B) comprend la cyclisation d'un composé de formule (VIII) :



(VIII)

dans laquelle Alk représente un groupe alkyle en C₁₋₆ et R¹ et R³ représentent ensemble une chaîne alkyle ou alcényle à 3 ou 4 chaînons, tels qu'ils sont tous deux définis plus haut; ou un procédé (C) pour préparer un composé de formule (I) dans laquelle R³ représente un groupe alkyle en C₁₋₃, lequel procédé (C) comprend la cyclisation d'un composé de formule (X) :




(X)


dans laquelle Alk représente un groupe alkyle en C₁₋₆ et R⁵ représente un groupe alkyle en C₂₋₅, substitué en C₁ par un atome d'halogène; ou un procédé (A), (B) ou (C) tel que décrit plus haut, suivi:

- i) d'une étape d'interconversion; et/ou
- ii) soit d'une formation de sel;
- iii) soit d'une formation de solvate.






16. Composés de formules (II), (III), (V), (VI), (VII), (VIII) et (X) :



(II)



(III)



30

où R^0 et R^2 sont tels que définis plus haut dans la revendication 1, R^1 et R^3 représentent ensemble une chaîne alkyle ou alcényle à 3 ou 4 chaînons, R^5 représente un groupe alkyle en C_{2-5} substitué en C_1 par un atome d'halogène, Alk représente un groupe alkyle en C_{1-6} et Hal est un atome d'halogène, à l'exception des composés de formules (III), (V), (VI) et (VII) dans lesquelles R^0 est un atome d'hydrogène, R^2 est un groupe phényle et Alk est un groupe méthyle.